

# Section 07: Induction & Midterm Review

---

## 1. Structural Induction

(a) Consider the following recursive definition of strings.

**Basis Step:** "" is a string

**Recursive Step:** If  $X$  is a string and  $c$  is a character then  $\text{append}(c, X)$  is a string.

Recall the following recursive definition of the function  $\text{len}$ :

$$\begin{aligned}\text{len}("") &= 0 \\ \text{len}(\text{append}(c, X)) &= 1 + \text{len}(X)\end{aligned}$$

Now, consider the following recursive definition:

$$\begin{aligned}\text{double}("") &= "" \\ \text{double}(\text{append}(c, X)) &= \text{append}(c, \text{append}(c, \text{double}(X))).\end{aligned}$$

Prove that for any string  $X$ ,  $\text{len}(\text{double}(X)) = 2\text{len}(X)$ .

(b) Consider the following definition of a (binary) **Tree**:

**Basis Step:**  $\bullet$  is a **Tree**.

**Recursive Step:** If  $L$  is a **Tree** and  $R$  is a **Tree** then  $\text{Tree}(\bullet, L, R)$  is a **Tree**.

The function  $\text{leaves}$  returns the number of leaves of a **Tree**. It is defined as follows:

$$\begin{aligned}\text{leaves}(\bullet) &= 1 \\ \text{leaves}(\text{Tree}(\bullet, L, R)) &= \text{leaves}(L) + \text{leaves}(R)\end{aligned}$$

Also, recall the definition of  $\text{size}$  on trees:

$$\begin{aligned}\text{size}(\bullet) &= 1 \\ \text{size}(\text{Tree}(\bullet, L, R)) &= 1 + \text{size}(L) + \text{size}(R)\end{aligned}$$

Prove that  $\text{leaves}(T) \geq \text{size}(T)/2 + 1/2$  for all **Trees**  $T$ .

(c) Prove the previous claim using strong induction. Define  $P(n)$  as "all trees  $T$  of size  $n$  satisfy  $\text{leaves}(T) \geq \text{size}(T)/2 + 1/2$ ". You may use the following facts:

- For any tree  $T$  we have  $\text{size}(T) \geq 1$ .
- For any tree  $T$ ,  $\text{size}(T) = 1$  if and only if  $T = \bullet$ .

If we wanted to prove these claims, we could do so by structural induction.

Note, in the inductive step you should start by letting  $T$  be an arbitrary tree of size  $k + 1$ .

## 2. Midterm Review: Translation

Let your domain of discourse be all coffee drinks. You should use the following predicates:

- $\text{soy}(x)$  is true iff  $x$  contains soy milk.

- $\text{whole}(x)$  is true iff  $x$  contains whole milk.
- $\text{sugar}(x)$  is true iff  $x$  contains sugar
- $\text{decaf}(x)$  is true iff  $x$  is not caffeinated.
- $\text{vegan}(x)$  is true iff  $x$  is vegan.
- $\text{RobbieLikes}(x)$  is true iff Robbie likes the drink  $x$ .

Translate each of the following statements into predicate logic. You may use quantifiers, the predicates above, and usual math connectors like  $=$  and  $\neq$ .

- Coffee drinks with whole milk are not vegan.
- Robbie only likes one coffee drink, and that drink is not vegan.
- There is a drink that has both sugar and soy milk.

Translate the following symbolic logic statement into a (natural) English sentence. Take advantage of domain restriction.

$$\forall x([\text{decaf}(x) \wedge \text{RobbieLikes}(x)] \rightarrow \text{sugar}(x))$$

### 3. Midterm Review: Number Theory

Let  $p$  be a prime number at least 3, and let  $x$  be an integer such that  $x^2 \not\equiv 1 \pmod{p}$ .

- Show that if an integer  $y$  satisfies  $y \equiv 1 \pmod{p}$ , then  $y^2 \equiv 1 \pmod{p}$ . (this proof will be short!)  
(Try to do this without using the theorem "Raising Congruences To A Power")
- Repeat part (a), but don't use any theorems from the Number Theory Reference Sheet. That is, show the claim directly from the definitions.
- From part (a), we can see that  $x \not\equiv 1 \pmod{p}$  can equal 1. Show that for any integer  $x$ , if  $x^2 \equiv 1 \pmod{p}$ , then  $x \equiv 1 \pmod{p}$  or  $x \equiv -1 \pmod{p}$ . That is, show that the only value  $x \not\equiv 1 \pmod{p}$  can take other than 1 is  $p - 1$ .  
Hint: Suppose you have an  $x$  such that  $x^2 \equiv 1 \pmod{p}$  and use the fact that  $x^2 - 1 = (x - 1)(x + 1)$   
Hint: You may use the following theorem without proof: if  $p$  is prime and  $p \mid (ab)$  then  $p \mid a$  or  $p \mid b$ .

### 4. Midterm Review: Induction

For any  $n \in \mathbb{N}$ , define  $S_n$  to be the sum of the squares of the first  $n$  positive integers, or

$$S_n = 1^2 + 2^2 + \cdots + n^2.$$

Prove that for all  $n \in \mathbb{N}$ ,  $S_n = \frac{1}{6}n(n+1)(2n+1)$ .