


Section 09: Solutions

1. Relations

- (a) Draw the transitive-reflexive closure of $\{(1, 2), (2, 3), (3, 4)\}$.

Solution:



- (b) Suppose that R is reflexive. Prove that $R \subseteq R^2$.

Solution:

Suppose $(a, b) \in R$. Since R is reflexive, we know $(b, b) \in R$ as well. Since there is a b such that $(a, b) \in R$ and $(b, b) \in R$, it follows that $(a, b) \in R^2$. Thus, $R \subseteq R^2$.

- (c) Consider the relation $R = \{(x, y) : x = y + 1\}$ on \mathbb{N} . Is R reflexive? Transitive? Symmetric? Anti-symmetric?

Solution:

It isn't reflexive, because $1 \neq 1 + 1$; so, $(1, 1) \notin R$. It isn't symmetric, because $(2, 1) \in R$ (because $2 = 1 + 1$), but $(1, 2) \notin R$, because $1 \neq 2 + 1$. It isn't transitive, because note that $(3, 2) \in R$ and $(2, 1) \in R$, but $(3, 1) \notin R$. It is anti-symmetric, because consider $(x, y) \in R$ such that $x \neq y$. Then, $x = y + 1$ by definition of R . However, $(y, x) \notin R$, because $y = x - 1 \neq x + 1$.

- (d) Consider the relation $S = \{(x, y) : x^2 = y^2\}$ on \mathbb{R} . Prove that S is reflexive, transitive, and symmetric.

Solution:

Consider $x \in \mathbb{R}$. Note that by definition of equality, $x^2 = x^2$; so, $(x, x) \in S$; so, S is reflexive.

Consider $(x, y) \in S$. Then, $x^2 = y^2$. It follows that $y^2 = x^2$; so, $(y, x) \in S$. So, S is symmetric.


Suppose $(x, y) \in S$ and $(y, z) \in S$. Then, $x^2 = y^2$, and $y^2 = z^2$. Since equality is transitive, $x^2 = z^2$. So, $(x, z) \in S$. So, S is transitive.

2. DFAs, Stage 1

Construct DFAs to recognize each of the following languages. Let $\Sigma = \{0, 1, 2, 3\}$.


- (a) All binary strings.

Solution:




- (b) All strings whose digits sum to an even number.

Solution:



- (c) All strings whose digits sum to an odd number.

Solution:




3. DFAs, Stage 2

Construct DFAs to recognize each of the following languages. Let $\Sigma = \{0, 1\}$.


- (a) All strings which do not contain the substring 101.

Solution:




q_1 : strings that don't contain 101 and end in 1.
 q_0 : $\varepsilon, 0$, strings that don't contain 101 and end in 00.

- (b) All strings containing at least two 0's and at most one 1.
Solution:




- (c) All strings containing an even number of 1's and an odd number of 0's and not containing the substring 10.
Solution:



4. NFAs

- (a) What language does the following NFA accept?




Solution:

All strings of only 0's and 1's not containing more than one 1.

(b) Create an NFA for the language “all binary strings that have a 1 as one of the last three digits”.

Solution:

The following is one such NFA:




5. DFAs & Minimization


Note: We will not test you on minimization, although you may optionally read the extra slides and do this problem for fun

(a) Convert the NFA from 1a to a DFA, then minimize it.


Solution:



Here is the minimized form:



(b) Minimize the following DFA:



Solution:

Step 1: q_0, q_2 are final states and the rest are not final. So, we start with the initial partition with the following groups: group 1 is $\{q_0, q_2\}$ and group 2 is $\{q_1, q_3, q_4\}$.

Step 2: q_1 is sending a to group 1 while q_3, q_4 are sending a to group 2. So, we divide group 2. We get the following groups: group 1 is $\{q_0, q_2\}$, group 3 is $\{q_1\}$ and group 4 is $\{q_3, q_4\}$.

Step 3: q_0 is sending a to group 3 and q_2 is sending a to group 4. So, we divide group 1. We will have the following groups: group 3 is $\{q_1\}$, group 4 is $\{q_3, q_4\}$, group 5 is $\{q_0\}$ and group 6 is $\{q_2\}$.

The minimized DFA is the following:

