

CSE 311 Section 6

Induction

Administrivia



Announcements & Reminders

- HW4
 - Grades out now
 - If you think something was graded incorrectly, submit a regrade request!
- HW5 (BOTH PARTS)
 - BOTH PARTS due Wednesday 10/8 @ 10pm
- Midterm is Coming!!!
 - Wednesday 10/15 @ 6-7:30 pm in BAG 131 and 154
 - If you cannot make it, please let us know ASAP and we will schedule you for a makeup

Induction



(Weak) Induction Template

Let $P(n)$ be “(whatever you’re trying to prove)”.

We show $P(n)$ holds for all n by induction on n .

Base Case: Show $P(b)$ is true.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq b$.

Inductive Step: Show $P(k + 1)$ (i.e. get $P(k) \rightarrow P(k + 1)$)

Conclusion: Therefore, $P(n)$ holds for all n by the principle of induction.

(Weak) Induction Template

Let $P(n)$ be “(whatever you’re trying to prove)”.
We show $P(n)$ holds **for all n** by induction on n .

Note: often you will
condition n here, like
“all natural numbers n ”
or “ $n \geq 0$ ”

Base Case: Show $P(b)$ is true.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq b$.

Inductive Step: Show $P(k + 1)$ (i.e. get $P(k) \rightarrow P(k + 1)$)

Conclusion: Therefore, $P(n)$ holds **for all n** by the principle of induction.

Match the earlier condition on n in your conclusion!

Problem 1 – Induction with Equality

- a) Show using induction that $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ for all $n \in \mathbb{N}$.
- b) Define the triangle numbers as $\Delta_n = 1 + 2 + \dots + n$, where $n \in \mathbb{N}$. In part (a) we showed $\Delta_n = \frac{n(n+1)}{2}$. Prove the following equality for all $n \in \mathbb{N}$:
- $$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Lets walk through part (a) together.

We can “fill in” our induction template to construct our proof by induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “”. We show $P(n)$ holds for (some) n by induction on n .

Base Case: $P(b)$:

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq b$.

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for (some) n by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(b)$:

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq b$.

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$.

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k + 1)$: $0 + 1 + \dots + k + (k + 1) = \frac{(k+1)(k+2)}{2}$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k + 1)$: $0 + 1 + \dots + k + (k + 1) = \frac{(k+1)(k+2)}{2}$

$$0 + 1 + \dots + k + (k + 1) = \dots$$

...

$$= \frac{(k+1)(k+2)}{2} \quad ?$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k + 1)$: $0 + 1 + \dots + k + (k + 1) = \frac{(k+1)(k+2)}{2}$

$$0 + 1 + \dots + k + (k + 1) = (0 + 1 + \dots + k) + (k + 1)$$

...

$$= \frac{(k+1)(k+2)}{2} \quad ?$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k + 1)$: $0 + 1 + \dots + k + (k + 1) = \frac{(k+1)(k+2)}{2}$

$$0 + 1 + \dots + k + (k + 1) = (0 + 1 + \dots + k) + (k + 1)$$

$$= \frac{k(k+1)}{2} + (k + 1) \quad \text{by I.H.}$$

...

$$= \frac{(k+1)(k+2)}{2} \quad ?$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k + 1)$: $0 + 1 + \dots + k + (k + 1) = \frac{(k+1)(k+2)}{2}$

$$0 + 1 + \dots + k + (k + 1) = (0 + 1 + \dots + k) + (k + 1)$$

$$= \frac{k(k+1)}{2} + (k + 1) \quad \text{by I.H.}$$

$$= \frac{k(k+1)}{2} + \frac{2(k+1)}{2}$$

...

$$= \frac{(k+1)(k+2)}{2} \quad ?$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k+1)$: $0 + 1 + \dots + k + (k+1) = \frac{(k+1)(k+2)}{2}$

$$\begin{aligned}0 + 1 + \dots + k + (k+1) &= (0 + 1 + \dots + k) + (k+1) \\ &= \frac{k(k+1)}{2} + (k+1) && \text{by I.H.} \\ &= \frac{k(k+1)}{2} + \frac{2(k+1)}{2} \\ &= \frac{k(k+1) + 2(k+1)}{2} \\ &= \frac{(k+1)(k+2)}{2} && ?\end{aligned}$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

Show using induction that
 $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$
for all $n \in \mathbb{N}$.

Let $P(n)$ be “ $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0 + \dots = 0 = \frac{0(0+1)}{2}$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$, i.e. $0 + 1 + 2 + \dots + k = \frac{k(k+1)}{2}$

Inductive Step: Goal: Show $P(k+1)$: $0 + 1 + \dots + k + (k+1) = \frac{(k+1)(k+2)}{2}$

$$\begin{aligned}0 + 1 + \dots + k + (k+1) &= (0 + 1 + \dots + k) + (k+1) \\ &= \frac{k(k+1)}{2} + (k+1) && \text{by I.H.} \\ &= \frac{k(k+1)}{2} + \frac{2(k+1)}{2} \\ &= \frac{k(k+1) + 2(k+1)}{2} \\ &= \frac{(k+1)(k+2)}{2} && \text{factoring out (k+1)}\end{aligned}$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

- a) Show using induction that $0 + 1 + 2 + \dots + n = \frac{n(n+1)}{2}$ for all $n \in \mathbb{N}$.
- b) Define the triangle numbers as $\Delta_n = 1 + 2 + \dots + n$, where $n \in \mathbb{N}$. In part (a) we showed $\Delta_n = \frac{n(n+1)}{2}$. Prove the following equality for all $n \in \mathbb{N}$:
- $$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Now try part (b) with people around you, and then we'll go over it together!

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “”. We show $P(n)$ holds for (some) n by induction on n .

Base Case: $P(b)$:

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq b$.

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for (some) n by the principle of induction.

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “ $0^3 + 1^3 + \dots + n^3 = (0 + 1 + \dots + n)^2$ ”. We show $P(n)$ holds for **all $n \in \mathbb{N}$** by induction on n .

Base Case: $P(b)$:

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq b$.

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for **all $n \in \mathbb{N}$** by the principle of induction.

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “ $0^3 + 1^3 + \dots + n^3 = (0 + 1 + \dots + n)^2$ ”. We show $P(n)$ holds for **all $n \in \mathbb{N}$** by induction on n .

Base Case: $P(0)$: $0^3 = 0 = (0)^2$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$.

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for **all $n \in \mathbb{N}$** by the principle of induction.

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “ $0^3 + 1^3 + \dots + n^3 = (0 + 1 + \dots + n)^2$ ”. We show $P(n)$ holds for **all** $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0^3 = 0 = (0)^2$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$. i.e. $0^3 + 1^3 + \dots + k^3 = (0 + 1 + \dots + k)^2$

Inductive Step: Goal: Show $P(k + 1)$:

Conclusion: Therefore, $P(n)$ holds for **all** $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “ $0^3 + 1^3 + \dots + n^3 = (0 + 1 + \dots + n)^2$ ”. We show $P(n)$ holds for **all** $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0^3 = 0 = (0)^2$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$. i.e. $0^3 + 1^3 + \dots + k^3 = (0 + 1 + \dots + k)^2$

Inductive Step: Goal: Show $P(k + 1)$: $0^3 + 1^3 + \dots + k^3 + (k + 1)^3 = (0 + 1 + \dots + k + (k + 1))^2$

Conclusion: Therefore, $P(n)$ holds for **all** $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “ $0^3 + 1^3 + \dots + n^3 = (0 + 1 + \dots + n)^2$ ”. We show $P(n)$ holds for **all** $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0^3 = 0 = (0)^2$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$. i.e. $0^3 + 1^3 + \dots + k^3 = (0 + 1 + \dots + k)^2$

Inductive Step: Goal: Show $P(k + 1)$: $0^3 + 1^3 + \dots + k^3 + (k + 1)^3 = (0 + 1 + \dots + k + (k + 1))^2$

$$0^3 + 1^3 + \dots + k^3 + (k + 1)^3 = \dots$$

...

$$= (0 + 1 + \dots + k + (k + 1))^2$$

Conclusion: Therefore, $P(n)$ holds for **all** $n \in \mathbb{N}$ by the principle of induction.

Problem 1 – Induction with Equality

$$\Delta_n = 1 + 2 + \dots + n, \quad n \in \mathbb{N}.$$

$$\Delta_n = \frac{n(n+1)}{2}. \quad \text{Prove for all } n \in \mathbb{N}:$$

$$0^3 + 1^3 + \dots + n^3 = \Delta_n^2$$

Let $P(n)$ be “ $0^3 + 1^3 + \dots + n^3 = (0 + 1 + \dots + n)^2$ ”. We show $P(n)$ holds for all $n \in \mathbb{N}$ by induction on n .

Base Case: $P(0)$: $0^3 = 0 = (0)^2$ so the base case holds.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 0$. i.e. $0^3 + 1^3 + \dots + k^3 = (0 + 1 + \dots + k)^2$

Inductive Step: Goal: Show $P(k + 1)$: $0^3 + 1^3 + \dots + k^3 + (k + 1)^3 = (0 + 1 + \dots + k + (k + 1))^2$

$$0^3 + 1^3 + \dots + k^3 + (k + 1)^3 = (0 + 1 + \dots + k)^2 + (k + 1)^3 \quad \text{by I.H.}$$

$$= \left(\frac{k(k+1)}{2}\right)^2 + (k + 1)^3 \quad \text{by (a)}$$

$$= (k + 1)^2 \left(\frac{k^2}{2^2} + (k + 1)\right) \quad \text{factor out } (k + 1)^2$$

$$= (k + 1)^2 \left(\frac{k^2 + 4k + 4}{4}\right)$$


$$= (k + 1)^2 \left(\frac{(k+2)^2}{4}\right) \quad \text{factor numerator}$$

$$= \left(\frac{(k+1)(k+2)}{2}\right)^2$$

$$= (0 + 1 + \dots + k + (k + 1))^2 \quad \text{by (a)}$$

Conclusion: Therefore, $P(n)$ holds for all $n \in \mathbb{N}$ by the principle of induction.

Strong Induction



Why Strong Induction?

In **weak induction**, the inductive hypothesis only assumes that $P(k)$ is true and uses that in the inductive step to prove the implication $P(k) \rightarrow P(k + 1)$.

In **strong induction**, the inductive hypothesis assumes the predicate holds for every step from the base case(s) up to $P(k)$. This usually looks something like $P(b_1) \wedge P(b_2) \wedge \dots \wedge P(k)$. Then it uses this stronger inductive hypothesis in the inductive step to prove the implication $P(b_1) \wedge \dots \wedge P(k) \rightarrow P(k + 1)$.

Strong induction is necessary when we have multiple base cases, or when we need to go back to a smaller number than k in our inductive step.

Strong Induction Template

Let $P(n)$ be “(whatever you’re trying to prove)”.

We show $P(n)$ holds for all $n \geq b_{min}$ by induction on n .

Base Case: Show $P(b_{min}), P(b_{min+1}), \dots, P(b_{max})$ are all true.

Inductive Hypothesis: Suppose $P(b_{min}) \wedge \dots \wedge P(k)$ hold for an arbitrary $k \geq b_{max}$.

Inductive Step: Show $P(k + 1)$ (i.e. get $P(b_{min}) \wedge \dots \wedge P(k) \rightarrow P(k + 1)$)

Conclusion: Therefore, $P(n)$ holds for all $n \geq b_{min}$ by the principle of induction.

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

First, let's construct a formula for $f(n)$. How many rabbits does he have each year? Let's do some calculations, and see if we can find a pattern. Then, we'll use induction to prove the pattern holds for all n !

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

$$f(0) = 0$$

$$f(1) = 1$$

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

$$f(0) = 0$$

$$f(1) = 1$$

$$f(2) = 2f(2 - 1) - f(2 - 2) = 2f(1) - f(0) = 2(1) - 0 = 2 - 0 = 2$$

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

$$f(0) = 0$$

$$f(1) = 1$$

$$f(2) = 2f(2 - 1) - f(2 - 2) = 2f(1) - f(0) = 2(1) - 0 = 2 - 0 = 2$$

$$f(3) = 2f(3 - 1) - f(3 - 2) = 2f(2) - f(1) = 2(2) - 1 = 4 - 1 = 3$$

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

$$f(0) = 0$$

$$f(1) = 1$$

$$f(2) = 2f(2 - 1) - f(2 - 2) = 2f(1) - f(0) = 2(1) - 0 = 2 - 0 = 2$$

$$f(3) = 2f(3 - 1) - f(3 - 2) = 2f(2) - f(1) = 2(2) - 1 = 4 - 1 = 3$$

$$f(4) = 2f(4 - 1) - f(4 - 2) = 2f(3) - f(2) = 2(3) - 2 = 6 - 2 = 4$$

Problem 4 – Cantelli's Rabbits

Xavier Cantelli owns some rabbits. The number of rabbits he has in any given year is described by the function f :

$$f(0) = 0$$

$$f(1) = 1$$

$$f(n) = 2f(n - 1) - f(n - 2) \text{ for } n \geq 2$$

Determine, with proof, the number, $f(n)$, of rabbits that Cantelli owns in year n . That is, construct a formula for $f(n)$ and prove its correctness.

$$f(0) = 0$$

$$f(1) = 1$$

$$f(2) = 2f(2 - 1) - f(2 - 2) = 2f(1) - f(0) = 2(1) - 0 = 2 - 0 = 2$$

$$f(3) = 2f(3 - 1) - f(3 - 2) = 2f(2) - f(1) = 2(2) - 1 = 4 - 1 = 3$$

$$f(4) = 2f(4 - 1) - f(4 - 2) = 2f(3) - f(2) = 2(3) - 2 = 6 - 2 = 4$$

It seems like we have a pattern here!

$$f(n) = n$$

But we don't want to have to check for EVERY n , so let's see if we can prove it with induction instead!

Problem 4 – Cantelli's Rabbits

What kind of induction should we use?

Problem 4 – Cantelli's Rabbits

What kind of induction should we use?

Strong induction!

Problem 4 – Cantelli's Rabbits

What kind of induction should we use?

Strong induction!

Two big clues:

- Multiple base cases in the formula: $f(0) = 0$ and $f(1) = 1$
- Recursively defined step of formula goes back further than just n :
 - $f(n)$ based on both $f(n - 1)$ and $f(n - 2)$
 - for $P(n)$ to be true, both $P(n - 1)$ and $P(n - 2)$ must be true

Problem 4 – Cantelli’s Rabbits

Let $P(n)$ be “(whatever you’re trying to prove)”.

We show $P(n)$ holds for all $n \geq b_{min}$ by induction on n .

Base Case: Show $P(b_{min}), P(b_{min+1}), \dots, P(b_{max})$ are all true.

Inductive Hypothesis: Suppose $P(b_{min}) \wedge \dots \wedge P(k)$ hold for an arbitrary $k \geq b_{max}$.

Inductive Step: Show $P(k + 1)$ (i.e. get $P(b_{min}) \wedge \dots \wedge P(k) \rightarrow P(k + 1)$)

Conclusion: Therefore, $P(n)$ holds for all $n \geq b_{min}$ by the principle of induction.

Fill in the strong induction template to prove the claim!

Problem 4 – Cantelli's Rabbits

Let $P(n)$ be “”.

We show $P(n)$ holds ...

Base Cases:

Inductive Hypothesis:

Inductive Step:

Conclusion: Therefore, $P(n)$ holds for all ... by the principle of induction.

Problem 4 – Cantelli's Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases:

Inductive Hypothesis:

Inductive Step:

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli's Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis:

Inductive Step:

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli’s Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis: Suppose $P(0) \wedge P(1) \wedge \cdots \wedge P(k)$ hold for an arbitrary all $k \geq 1$.

Inductive Step:

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli's Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis: Suppose $P(0) \wedge P(1) \wedge \cdots \wedge P(k)$ hold for an arbitrary all $k \geq 1$.
i.e. $f(k) = k, f(k - 1) = k - 1, f(k - 2) = k - 2$, etc.

Inductive Step:

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli's Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis: Suppose $P(0) \wedge P(1) \wedge \cdots \wedge P(k)$ hold for an arbitrary all $k \geq 1$.

i.e. $f(k) = k, f(k - 1) = k - 1, f(k - 2) = k - 2$, etc.

Inductive Step: Goal: Show $P(k + 1): f(k + 1) = k + 1$

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli’s Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis: Suppose $P(0) \wedge P(1) \wedge \dots \wedge P(k)$ hold for an arbitrary all $k \geq 1$.

i.e. $f(k) = k, f(k - 1) = k - 1, f(k - 2) = k - 2$, etc.

Inductive Step: Goal: Show $P(k + 1): f(k + 1) = k + 1$

$$f(k + 1) = \dots$$

...

$$= k + 1$$

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli’s Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis: Suppose $P(0) \wedge P(1) \wedge \dots \wedge P(k)$ hold for an arbitrary all $k \geq 1$.

i.e. $f(k) = k, f(k - 1) = k - 1, f(k - 2) = k - 2$, etc.

Inductive Step: Goal: Show $P(k + 1): f(k + 1) = k + 1$

$$f(k + 1) = 2f(k) - f(k - 1) \quad \text{definition of } f$$

...

$$= k + 1$$

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

Problem 4 – Cantelli’s Rabbits

Let $P(n)$ be “ $f(n) = n$ ”.

We show $P(n)$ holds for all $n \geq 0$ by induction on n .

Base Cases: ($n = 0, n = 1$): $f(0) = 0$ and $f(1) = 1$ by definition of f .

Inductive Hypothesis: Suppose $P(0) \wedge P(1) \wedge \dots \wedge P(k)$ hold for an arbitrary all $k \geq 1$.

i.e. $f(k) = k, f(k - 1) = k - 1, f(k - 2) = k - 2$, etc.

Inductive Step: Goal: Show $P(k + 1)$: $f(k + 1) = k + 1$

$$\begin{aligned} f(k + 1) &= 2f(k) - f(k - 1) && \text{definition of } f \\ &= 2(k) - (k - 1) && \text{by I.H.} \\ &= k + 1 \end{aligned}$$

Conclusion: Therefore, $P(n)$ holds for all $n \geq 0$ by the principle of induction.

That's All, Folks!

Thanks for coming to section this week!
Any questions?