

Warm up:


What is the following recursively-defined set?

Basis Step: $4 \in S, 5 \in S$

Recursive Step: If $x \in S$ and $y \in S$ then $x - y \in S$

Structural Induction and Regular Expressions

CSE 311 Winter 2024
Lecture 17



Trees!




More Structural Sets

Binary Trees are another common source of structural induction.

Basis: A single node is a rooted binary tree. ●

Recursive Step: If T_1 and T_2 are rooted binary trees with roots r_1 and r_2 , then a tree rooted at a new node, with children r_1, r_2 is a binary tree.



Functions on Binary Trees

$$\text{size}(\bullet) = 1$$

$$\text{size}\left(\begin{array}{c} \bullet \\ / \quad \backslash \\ \bullet \quad \bullet \\ \triangleleft \quad \triangleright \\ T_1 \quad T_2 \end{array}\right) = \text{size}(T_1) + \text{size}(T_2) + 1$$

$$\text{height}(\bullet) = 0$$

$$\text{height}\left(\begin{array}{c} \bullet \\ / \quad \backslash \\ \bullet \quad \bullet \\ \triangleleft \quad \triangleright \\ T_1 \quad T_2 \end{array}\right) = 1 + \max(\text{height}(T_1), \text{height}(T_2))$$

Claim

We want to show that trees of a certain height can't have too many nodes. Specifically our claim is this:

For all trees T , $\text{size}(T) \leq 2^{\text{height}(T)+1} - 1$


Take a moment to absorb this formula, then we'll do induction!

Structural Induction on Binary Trees

Let $P(T)$ be " $\text{size}(T) \leq 2^{\text{height}(T)+1} - 1$ ". We show $P(T)$ for all binary trees T by structural induction.

Base Case: Let $T = \bullet$. $\text{size}(T)=1$ and $\text{height}(T) = 0$, so $\text{size}(T)=1 \leq 2 - 1 = 2^{0+1} - 1 = 2^{\text{height}(T)+1} - 1$.


Inductive Hypothesis: Suppose $P(L)$ and $P(R)$ hold for arbitrary trees L, R . Let T be the tree



Inductive step: Figure out, (1) what we must show (2) a formula for height and a formula for size of T .

Structural Induction on Binary Trees (cont.)

Let $P(T)$ be " $\text{size}(T) \leq 2^{\text{height}(T)+1} - 1$ ". We show $P(T)$ for all binary trees T by structural induction.




$$\text{height}(T) = 1 + \max\{\text{height}(L), \text{height}(R)\}$$

$$\text{size}(T) = 1 + \text{size}(L) + \text{size}(R)$$


So $P(T)$ holds, and we have $P(T)$ for all binary trees T by the principle of induction.

How do heights compare?


If L is taller than R ?



If L, R same height?




If R is taller than L ?



$$\text{height}(\bullet) = 0$$


$$\text{height}(\text{tree}) =$$



$$1 + \max(\text{height}(T_1), \text{height}(T_2))$$

How do heights compare?


If L is taller than R ?



$$\text{height}(T) = \text{height}(L) + 1$$

$$\text{height}(T) > \text{height}(R) + 1$$


If L, R same height?



$$\text{height}(T) = \text{height}(L) + 1$$

$$\text{height}(T) = \text{height}(R) + 1$$

If R is taller than L ?




$$\text{height}(T) > \text{height}(L) + 1$$

$$\text{height}(T) = \text{height}(R) + 1$$

In all cases: $\text{height}(T) \geq \text{height}(L) + 1$, $\text{height}(T) \geq \text{height}(R) + 1$

Structural Induction on Binary Trees (cont.)

Let $P(T)$ be " $\text{size}(T) \leq 2^{\text{height}(T)+1} - 1$ ". We show $P(T)$ for all binary trees T by structural induction.



$$\text{height}(T) = 1 + \max\{\text{height}(L), \text{height}(R)\}$$

$$\text{size}(T) = 1 + \text{size}(L) + \text{size}(R)$$

$$\text{size}(T) = 1 + \text{size}(L) + \text{size}(R) \leq 1 + 2^{\text{height}(L)+1} - 1 + 2^{\text{height}(R)+1} - 1 \quad (\text{by IH})$$


$$\leq 2^{\text{height}(L)+1} + 2^{\text{height}(R)+1} - 1 \quad (\text{cancel 1's})$$

$$\leq 2^{\text{height}(T)} + 2^{\text{height}(T)} - 1 = 2^{\text{height}(T)+1} - 1 \quad (T \text{ taller than subtrees})$$

So $P(T)$ holds, and we have $P(T)$ for all binary trees T by the principle of induction.

Structural Induction Template

1. Define $P()$ State that you will show $P(x)$ holds for all $x \in S$ and that your proof is by structural induction.
2. Base Case: Show $P(b)$
[Do that for every b in the basis step of defining S]
3. Inductive Hypothesis: Suppose $P(x)$
[Do that for every x listed as already in S in the recursive rules].
4. Inductive Step: Show $P()$ holds for the "new elements."
[You will need a separate step for every element created by the recursive rules].
5. Therefore $P(x)$ holds for all $x \in S$ by the principle of induction.



Structural Induction on Strings

Strings

ε is "the empty string"

The string with 0 characters – "" in Java (not null!)

Σ^* :

Basis: $\varepsilon \in \Sigma^*$.

Recursive: If $w \in \Sigma^*$ and $a \in \Sigma$ then $wa \in \Sigma^*$

wa means the string of w with the character a appended.

You'll also see $w \cdot a$ ($a \cdot$ to mean "concatenate" i.e. + in Java)

Functions on Strings

Since strings are defined recursively, most functions on strings are as well.

Length:

$$\text{len}(\varepsilon) = 0;$$

$$\text{len}(wa) = \text{len}(w) + 1 \text{ for } w \in \Sigma^*, a \in \Sigma$$

Reversal:

$$\varepsilon^R = \varepsilon;$$

$$(wa)^R = aw^R \text{ for } w \in \Sigma^*, a \in \Sigma$$

Concatenation

$$x \cdot \varepsilon = x \text{ for all } x \in \Sigma^*;$$

$$x \cdot (wa) = (x \cdot w)a \text{ for } w \in \Sigma^*, a \in \Sigma$$

Number of c 's in a string

$$\#_c(\varepsilon) = 0$$

$$\#_c(wc) = \#_c(w) + 1 \text{ for } w \in \Sigma^*;$$

$$\#_c(wa) = \#_c(w) \text{ for } w \in \Sigma^*, a \in \Sigma \setminus \{c\}.$$

Claim for all $x, y \in \Sigma^*$ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$.

Let $P(y)$ be "for all $x \in \Sigma^*$ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$."

Notice the strangeness of this $P()$ there is a "for all x " inside the definition of $P(y)$.

That means we'll have to introduce an arbitrary x as part of the base case and the inductive step!

Claim for all $x, y \in \Sigma^*$ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$.

Let $P(y)$ be " $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$ for all $x \in \Sigma^*$."

We prove $P(y)$ for all $x \in \Sigma^*$ by structural induction.

Base Case:

Inductive Hypothesis

Inductive Step:

We conclude that $P(y)$ holds for all string y by the principle of induction. Unwrapping the definition of P , we get $\forall x \forall y \in \Sigma^* \text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

Claim for all $x, y \in \Sigma^*$ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$.

Let $P(y)$ be " $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$ for all $x \in \Sigma^*$."

We prove $P(y)$ for all $x \in \Sigma^*$ by structural induction.

Base Case: Let x be an arbitrary string, $\text{len}(x \cdot \epsilon) = \text{len}(x)$
 $= \text{len}(x) + 0 = \text{len}(x) + \text{len}(\epsilon)$

Inductive Hypothesis: Suppose $P(w)$ for an arbitrary string w .

Inductive Step:

We conclude that $P(y)$ holds for all string y by the principle of induction. Unwrapping the definition of P , we get $\forall x \forall y \in \Sigma^* \text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

Claim for all $x, y \in \Sigma^*$ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$.

Let $P(y)$ be " $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$ for all $x \in \Sigma^*$."

We prove $P(y)$ for all $x \in \Sigma^*$ by structural induction.

Base Case: Let x be an arbitrary string, $\text{len}(x \cdot \epsilon) = \text{len}(x)$
 $= \text{len}(x) + 0 = \text{len}(x) + \text{len}(\epsilon)$

Inductive Hypothesis: Suppose $P(w)$ for an arbitrary string w .

Inductive Step: Let $y = wa$ for an arbitrary $a \in \Sigma$. We show $P(y)$. Let x be an arbitrary string.

...

Therefore, $\text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

We conclude that $P(y)$ holds for all string y by the principle of induction. Unwrapping the definition of P , we get $\forall x \forall y \in \Sigma^* \text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

Claim for all $x, y \in \Sigma^*$ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$.

Let $P(y)$ be “ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$ for all $x \in \Sigma^*$.”

We prove $P(y)$ for all $x \in \Sigma^*$ by structural induction.

Base Case: Let x be an arbitrary string, $\text{len}(x \cdot \epsilon) = \text{len}(x)$
 $= \text{len}(x) + 0 = \text{len}(x) + \text{len}(\epsilon)$

Inductive Hypothesis: Suppose $P(w)$ for an arbitrary string w .

Inductive Step: Let $y = wa$ for an arbitrary $a \in \Sigma$. We show $P(y)$. Let x be an arbitrary string.

$\text{len}(xy) = \text{len}(xwa) = \text{len}(xw) + 1$ (by definition of len)

$= \text{len}(x) + \text{len}(w) + 1$ (by IH)

$= \text{len}(x) + \text{len}(wa)$ (by definition of len)

Therefore, $\text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

We conclude that $P(y)$ holds for all string y by the principle of induction. Unwrapping the definition of P , we get $\forall x \forall y \in \Sigma^* \text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

Why all those arbitraries?

Let $P(y)$ be “ $\text{len}(x \cdot y) = \text{len}(x) + \text{len}(y)$ for all $x \in \Sigma^*$.”

We prove $P(y)$ for all $x \in \Sigma^*$ by structural induction.

Base Case: Let x be an arbitrary string, $\text{len}(x \cdot \epsilon) = \text{len}(x) = \text{len}(x) + 0 = \text{len}(x) + \text{len}(\epsilon)$

Inductive Hypothesis: Suppose $P(w)$ for an arbitrary string w .

Inductive Step: Let $y = wa$ for an arbitrary $a \in \Sigma$. We show $P(y)$. Let x be an arbitrary string.

$\text{len}(xy) = \text{len}(xwa) = \text{len}(xw) + 1$ (by definition of len)

$= \text{len}(x) + \text{len}(w) + 1$ (by IH)

$= \text{len}(x) + \text{len}(wa)$ (by definition of len)

Therefore, $\text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.


We conclude that $P(y)$ holds for all strings y by the principle of induction. Unwrapping the definition of P , we get $\forall x \forall y \in \Sigma^* \text{len}(xy) = \text{len}(x) + \text{len}(y)$, as required.

$P(\epsilon)$ is a for-all statement, introduce arbitrary variable to show for-all.

Needs to be arbitrary because it's in the IH (induction wouldn't show “all strings” otherwise)

Recursive rule says “every $a \in \Sigma$ ” so we need to argue for every a .

$P(y)$ is a for-all statement, introduce arbitrary variable to show for-all.



A few last comments

What does the inductive step look like?

Here's a recursively-defined set:

Basis: $0 \in T$ and $5 \in T$

Recursive: If $x, y \in T$ then $x + y \in T$ and $x - y \in T$.

Let $P(x)$ be " $5|x$ "

What does the inductive step look like?

Well there's two recursive rules, so we have two things to show

Just the IS (you still need the other steps)

Let t be an arbitrary element of T not covered by the base case. By the exclusion rule $t = x + y$ or $t = x - y$ for $x, y \in T$.

Inductive hypothesis: Suppose $P(x)$ and $P(y)$ hold.

Case 1: $t = x + y$

By IH $5|x$ and $5|y$ so $5a = x$ and $5b = y$ for integers a, b .

Adding, we get $x + y = 5a + 5b = 5(a + b)$. Since a, b are integers, so is $a + b$, and $P(x + y)$, i.e. $P(t)$, holds.

Case 2: $t = x - y$

By IH $5|x$ and $5|y$ so $5a = x$ and $5b = y$ for integers a, b .

Subtracting, we get $x - y = 5a - 5b = 5(a - b)$. Since a, b are integers, so is $a - b$, and $P(x - y)$, i.e., $P(t)$, holds.

In all cases, we have $P(t)$. By the principle of induction, $P(x)$ holds for all $x \in T$.

If you don't have a recursively-defined set


You won't do structural induction.

You can do weak or strong induction though.

For example, Let $P(n)$ be "for all elements of S of "size" n <something> is true"

To prove "for all $x \in S$ of size n ..." you need to start with "let x be an arbitrary element of size $k + 1$ in your IS.

You CAN'T start with size k and "build up" to an arbitrary element of size $k + 1$ it isn't arbitrary.



Part 3 of the course!

Course Outline

Symbolic Logic (training wheels)

Just make arguments in mechanical ways.

Set Theory/Number Theory (bike in your backyard)

Models of computation (biking in your neighborhood)

Still make and communicate rigorous arguments

But now with objects you haven't used before.

- A first taste of how we can argue rigorously about computers.

First up: regular expressions, context free grammars, automata – understand these “simpler computers”

Soon: what these simple computers can do

Then: what simple computers can't do.

Last week: A problem our computers cannot solve.



The definitions for Friday



Regular Expressions

I have a giant text document. And I want to find all the email addresses inside. What does an email address look like?

[some letters and numbers] @ [more letters] . [com, net, or edu]

We want to ctrl-f for a **pattern of strings** rather than a single string

Languages

A set of strings is called a **language**.

Σ^* is a language

“the set of all binary strings of even length” is a language.

“the set of all palindromes” is a language.

“the set of all English words” is a language.

“the set of all strings matching a given **pattern**” is a language.

Regular Expressions

Basis:

ε is a regular expression. The empty string itself matches the pattern (and nothing else does).

\emptyset is a regular expression. No strings match this pattern.

a is a regular expression, for any $a \in \Sigma$ (i.e. any character). The character itself matching this pattern.

Recursive

If A, B are regular expressions then $(A \cup B)$ is a regular expression
matched by any string that matches A or that matches B [or both]).

If A, B are regular expressions then AB is a regular expression.
matched by any string x such that $x = yz$, y matches A and z matches B .

If A is a regular expression, then A^* is a regular expression.
matched by any string that can be divided into 0 or more strings that match A .


Regular Expressions

$(a \cup bc)$

$0(0 \cup 1)1$

0^*

$(0 \cup 1)^*$



Extra Practice



Induction: Hats!

You have n people in a line ($n \geq 2$). Each of them wears either a **purple hat** or a **gold hat**. The person at the front of the line wears a purple hat. The person at the back of the line wears a gold hat.

Show that for every arrangement of the line satisfying the rule above, there is a person with a purple hat next to someone with a gold hat.

Yes, this is kinda obvious. I promise this is good induction practice.

Yes, you could argue this by contradiction. I promise this is good induction practice.

Induction: Hats!

Define $P(n)$ to be "in every line of n people with gold and purple hats, with a purple hat at one end and a gold hat at the other, there is a person with a purple hat next to someone with a gold hat"

We show $P(n)$ for all integers $n \geq 2$ by induction on n .

Base Case: $n = 2$

Inductive Hypothesis:

Inductive Step:

By the principle of induction, we have $P(n)$ for all $n \geq 2$

Induction: Hats!

Define $P(n)$ to be "in every line of n people with gold and purple hats, with a purple hat at one end and a gold hat at the other, there is a person with a purple hat next to someone with a gold hat"

We show $P(n)$ for all integers $n \geq 2$ by induction on n .

Base Case: $n = 2$ The line must be just a person with a purple hat and a person with a gold hat, who are next to each other.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 2$.

Inductive Step: Consider an arbitrary line with $k + 1$ people in purple and gold hats, with a gold hat at one end and a purple hat at the other.

Target: there is someone in a purple hat next to someone in a gold hat.

By the principle of induction, we have $P(n)$ for all $n \geq 2$

Induction: Hats!

Define $P(n)$ to be "in every line of n people with gold and purple hats, with a purple hat at one end and a gold hat at the other, there is a person with a purple hat next to someone with a gold hat"

We show $P(n)$ for all integers $n \geq 2$ by induction on n .

Base Case: $n = 2$ The line must be just a person with a purple hat and a person with a gold hat, who are next to each other.

Inductive Hypothesis: Suppose $P(k)$ holds for an arbitrary $k \geq 2$.

Inductive Step: Consider an arbitrary line with $k + 1$ people in purple and gold hats, with a gold hat at one end and a purple hat at the other.

Case 1: There is someone with a purple hat next to the person in the gold hat at one end. Then those people are the required adjacent opposite hats.

Case 2: There is a person with a gold hat next to the person in the gold hat at the end. Then the line from the second person to the end is length k , has a gold hat at one end and a purple hat at the other. Applying the inductive hypothesis, there is an adjacent, opposite-hat wearing pair.

In either case we have $P(k + 1)$.

By the principle of induction, we have $P(n)$ for all $n \geq 2$