

CSE 312

Foundations of Computing II

Lecture 17: CLT and Polling



Anna R. Karlin

Slide Credit: Based on Stefano Tessaro's slides for 312 19au incorporating ideas from Alex Tsun, Rachel Lin, Hunter Schafer & myself 😊

The Normal Distribution

Definition. A **Gaussian (or normal) random variable** with parameters $\mu \in \mathbb{R}$ and $\sigma \geq 0$ has density

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$


(We say that X follows the Normal Distribution, and write $X \sim \mathcal{N}(\mu, \sigma^2)$)

Fact. If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $\mathbb{E}(X) = \mu$, and $\text{Var}(X) = \sigma^2$

Proof is easy because density curve is symmetric around μ , $f_X(\mu - x) = f_X(\mu + x)$

The Normal Distribution

Aka a “Bell Curve” (imprecise name)



CDF of normal distribution

Fact. If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Y = aX + b \sim \mathcal{N}(a\mu + b, a^2\sigma^2)$

Standard (unit) normal $Z \sim \mathcal{N}(0, 1)$


CDF. $\Phi(z) = \mathbb{P}(Z \leq z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^z e^{-x^2/2} dx$ for $Z \sim \mathcal{N}(0, 1)$

Note: $\Phi(z)$ has no closed form – generally given via tables

Table of $\Phi(z)$ CDF of Standard Normal Distn

Φ Table: $\mathbb{P}(Z \leq z)$ when $Z \sim \mathcal{N}(0, 1)$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.5279	0.53188	0.53586
0.1	0.53983	0.5438	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535
0.2	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409
0.3	0.61791	0.62172	0.62552	0.6293	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173
0.4	0.65542	0.6591	0.66276	0.6664	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793
0.5	0.69146	0.69497	0.69847	0.70194	0.7054	0.70884	0.71226	0.71566	0.71904	0.7224
0.6	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.7549
0.7	0.75804	0.76115	0.76424	0.7673	0.77035	0.77337	0.77637	0.77935	0.7823	0.78524
0.8	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327
0.9	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891
1.0	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214
1.1	0.86433	0.8665	0.86864	0.87076	0.87286	0.87493	0.87698	0.879	0.881	0.88298
1.2	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147
1.3	0.9032	0.9049	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774
1.4	0.91924	0.92073	0.9222	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189
1.5	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408
1.6	0.9452	0.9463	0.94738	0.94845	0.9495	0.95053	0.95154	0.95254	0.95352	0.95449
1.7	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.9608	0.96164	0.96246	0.96327
1.8	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062
1.9	0.97128	0.97193	0.97257	0.9732	0.97381	0.97441	0.975	0.97558	0.97615	0.9767
2.0	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.9803	0.98077	0.98124	0.98169
2.1	0.98214	0.98257	0.983	0.98341	0.98382	0.98422	0.98461	0.985	0.98537	0.98574
2.2	0.9861	0.98645	0.98679	0.98713	0.98745	0.98778	0.98809	0.9884	0.9887	0.98899
2.3	0.98928	0.98956	0.98983	0.9901	0.99036	0.99061	0.99086	0.99111	0.99134	0.99158
2.4	0.9918	0.99202	0.99224	0.99245	0.99266	0.99286	0.99305	0.99324	0.99343	0.99361
2.5	0.99379	0.99396	0.99413	0.9943	0.99446	0.99461	0.99477	0.99492	0.99506	0.9952
2.6	0.99534	0.99547	0.9956	0.99573	0.99585	0.99598	0.99609	0.99621	0.99632	0.99643
2.7	0.99653	0.99664	0.99674	0.99683	0.99693	0.99702	0.99711	0.9972	0.99728	0.99736
2.8	0.99744	0.99752	0.9976	0.99767	0.99774	0.99781	0.99788	0.99795	0.99801	0.99807
2.9	0.99813	0.99819	0.99825	0.99831	0.99836	0.99841	0.99846	0.99851	0.99856	0.99861
3.0	0.99865	0.99869	0.99874	0.99878	0.99882	0.99886	0.99889	0.99893	0.99896	0.999



What about Non-standard normal?

If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $\frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$

Therefore,


$$F_X(z) = \mathbb{P}(X \leq z) = \mathbb{P}\left(\frac{X - \mu}{\sigma} \leq \frac{z - \mu}{\sigma}\right) = \Phi\left(\frac{z - \mu}{\sigma}\right)$$

Agenda


- Central Limit Theorem (CLT) ◀
- Polling

CLT → empirical distribution of data often Gaussian


Neuron Activity



Vegetables



S&P 500 Returns after Elections



Examples from:
<https://galtonboard.com/probabilityexamplesinlife>

Sum of Independent RVs

i.i.d. = independent and identically distributed

X_1, \dots, X_n i.i.d. with expectation μ and variance σ^2

Define

$$S_n = X_1 + \dots + X_n$$

$$\mathbb{E}(S_n) = \mathbb{E}(X_1) + \dots + \mathbb{E}(X_n) = n\mu$$

$$\text{Var}(S_n) = \text{Var}(X_1) + \dots + \text{Var}(X_n) = n\sigma^2$$

Empirical observation: S_n looks like a normal RV as n grows.

Setup for Central Limit Theorem

X_1, \dots, X_n i.i.d., each with expectation μ and variance σ^2

Define $S_n = X_1 + \dots + X_n$ and

$$Y_n = \frac{S_n - n\mu}{\sigma\sqrt{n}}$$

$$\mathbb{E}(Y_n) = \frac{1}{\sigma\sqrt{n}} (\mathbb{E}(S_n) - n\mu) = \frac{1}{\sigma\sqrt{n}} (n\mu - n\mu) = 0$$

$$\text{Var}(Y_n) = \frac{1}{\sigma^2 n} (\text{Var}(S_n - n\mu)) = \frac{\text{Var}(S_n)}{\sigma^2 n} = \frac{\sigma^2 n}{\sigma^2 n} = 1$$

Central Limit Theorem

$$Y_n = \frac{X_1 + \cdots + X_n - n\mu}{\sigma\sqrt{n}}$$

Theorem. (Central Limit Theorem) The CDF of Y_n converges to the CDF of the standard normal $\mathcal{N}(0,1)$, i.e.,

$$\lim_{n \rightarrow \infty} \mathbb{P}(Y_n \leq y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^y e^{-x^2/2} dx$$

Also stated as:

- $\lim_{n \rightarrow \infty} Y_n \rightarrow \mathcal{N}(0,1)$
- $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n X_i \rightarrow \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$ where $\mu = E[X_i]$ and $\sigma^2 = \text{Var}(X_i)$

Agenda

- Central Limit Theorem (CLT)
- Polling ◀

Magic Mushrooms

Not that long ago, Oregonians voted on whether to legalize the therapeutic use of “magic mushrooms”.

Poll to determine the fraction of the population that will vote in favor of legalization.

- Call up a random sample of n people to ask their opinion
- Report the empirical fraction

Questions

- Is this a good estimate?
- How to choose n ?



Polling Accuracy

Often see claims that say

“Our poll found 80% support. This poll is accurate to within 5% with 98% probability”

Will unpack what this means and how they sample enough people to know this is true.

Formalizing Polls

Population size N , true fraction of voting in favor p , sample size n .

Problem: We don't know p

Polling Procedure

for $i = 1 \dots n$:

1. Pick uniformly random person to call (prob: $1/N$)
2. Ask them how they will vote

$$X_i = \begin{cases} 1, & \text{voting in favor} \\ 0, & \text{otherwise} \end{cases}$$

Report our estimate of p :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Formalizing Polls

Population size N , true fraction of voting in favor p , sample size n .

Problem: We don't know p

Polling Procedure

for $i = 1 \dots n$:

1. Pick uniformly random person to call (prob: $1/N$)
2. Ask them how they will vote

$$X_i = \begin{cases} 1, & \text{voting in favor} \\ 0, & \text{otherwise} \end{cases}$$

Report our estimate of p :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$$

Poll what type of r.v. is X_i ?

	Type	$E[X_i]$	$Var(X_i)$
a.	Bernoulli	p	$p(1-p)$
b.	Bernoulli	p	p^2
c.	Geometric	p	$\frac{1-p}{p^2}$
d.	Binomial	np	$np(1-p)$

<https://pollev.com/annakarlin185>

Random Variables

What type of r.v. is X_i ?

What can you say about

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i?$$

<https://pollev.com/annakarlin185>

Poll:

	$E[\bar{X}]$	$Var(\bar{X})$
a.	np	$np(1-p)$
b.	p	$p(1-p)$
c.	p	$p(1-p)/n$
d.	p/n	$p(1-p)/n$

Roadmap: Bounding Error

Goal: Find the value of n such that 98% of the time, the estimate \bar{X} is within 5% of the true p .

Central Limit Theorem

With i.i.d random variables X_1, X_2, \dots, X_n where
 $E[X_i] = \mu$ and $Var(X_i) = \sigma^2$

As $n \rightarrow \infty$,

$$\frac{X_1 + X_2 + \dots + X_n - n\mu}{\sigma\sqrt{n}} \rightarrow \mathcal{N}(0, 1)$$

Restated: As $n \rightarrow \infty$,

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \rightarrow \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$$

Central Limit Theorem

With i.i.d random variables X_1, X_2, \dots, X_n where $E[X_i] = \mu$ and $Var(X_i) = \sigma^2$

As $n \rightarrow \infty$,

$$\frac{X_1 + X_2 + \dots + X_n - n\mu}{\sigma\sqrt{n}} \rightarrow \mathcal{N}(0, 1)$$

Restated: As $n \rightarrow \infty$,

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \rightarrow \mathcal{N}\left(\mu, \frac{\sigma^2}{n}\right)$$

Poll: In the limit \bar{X} is ?

- a. $\mathcal{N}(0, 1)$
- b. $\mathcal{N}(p, p(1 - p))$
- c. $\mathcal{N}(p, p(1 - p)/n)$
- d. I don't know.

Roadmap: Bounding Error

Goal: Find the value of n such that 98% of the time, the estimate \bar{X} is within 5% of the true p

1. Define probability of a “bad event”
2. Apply CLT
3. Convert to a standard normal
4. Solve for n


Define probability of a “bad event”

Apply CLT

Convert to a standard normal

Solve for n

Table of $\Phi(z)$ CDF of Standard Normal Distn



Φ Table: $\mathbb{P}(Z \leq z)$ when $Z \sim \mathcal{N}(0, 1)$

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5	0.50399	0.50798	0.51197	0.51595	0.51994	0.52392	0.5279	0.53188	0.53586
0.1	0.53983	0.5438	0.54776	0.55172	0.55567	0.55962	0.56356	0.56749	0.57142	0.57535
0.2	0.57926	0.58317	0.58706	0.59095	0.59483	0.59871	0.60257	0.60642	0.61026	0.61409
0.3	0.61791	0.62172	0.62552	0.6293	0.63307	0.63683	0.64058	0.64431	0.64803	0.65173
0.4	0.65542	0.6591	0.66276	0.6664	0.67003	0.67364	0.67724	0.68082	0.68439	0.68793
0.5	0.69146	0.69497	0.69847	0.70194	0.7054	0.70884	0.71226	0.71566	0.71904	0.7224
0.6	0.72575	0.72907	0.73237	0.73565	0.73891	0.74215	0.74537	0.74857	0.75175	0.7549
0.7	0.75804	0.76115	0.76424	0.7673	0.77035	0.77337	0.77637	0.77935	0.7823	0.78524
0.8	0.78814	0.79103	0.79389	0.79673	0.79955	0.80234	0.80511	0.80785	0.81057	0.81327
0.9	0.81594	0.81859	0.82121	0.82381	0.82639	0.82894	0.83147	0.83398	0.83646	0.83891
1.0	0.84134	0.84375	0.84614	0.84849	0.85083	0.85314	0.85543	0.85769	0.85993	0.86214
1.1	0.86433	0.8665	0.86864	0.87076	0.87286	0.87493	0.87698	0.879	0.881	0.88298
1.2	0.88493	0.88686	0.88877	0.89065	0.89251	0.89435	0.89617	0.89796	0.89973	0.90147
1.3	0.9032	0.9049	0.90658	0.90824	0.90988	0.91149	0.91309	0.91466	0.91621	0.91774
1.4	0.91924	0.92073	0.9222	0.92364	0.92507	0.92647	0.92785	0.92922	0.93056	0.93189
1.5	0.93319	0.93448	0.93574	0.93699	0.93822	0.93943	0.94062	0.94179	0.94295	0.94408
1.6	0.9452	0.9463	0.94738	0.94845	0.9495	0.95053	0.95154	0.95254	0.95352	0.95449
1.7	0.95543	0.95637	0.95728	0.95818	0.95907	0.95994	0.9608	0.96164	0.96246	0.96327
1.8	0.96407	0.96485	0.96562	0.96638	0.96712	0.96784	0.96856	0.96926	0.96995	0.97062
1.9	0.97128	0.97193	0.97257	0.9732	0.97381	0.97441	0.975	0.97558	0.97615	0.9767
2.0	0.97725	0.97778	0.97831	0.97882	0.97932	0.97982	0.9803	0.98077	0.98124	0.98169
2.1	0.98214	0.98257	0.983	0.98341	0.98382	0.98422	0.98461	0.985	0.98537	0.98574
2.2	0.9861	0.98645	0.98679	0.98713	0.98745	0.98778	0.98809	0.9884	0.9887	0.98899
2.3	0.98928	0.98956	0.98983	0.9901	0.99036	0.99061	0.99086	0.99111	0.99134	0.99158
2.4	0.9918	0.99202	0.99224	0.99245	0.99266	0.99286	0.99305	0.99324	0.99343	0.99361
2.5	0.99379	0.99396	0.99413	0.9943	0.99446	0.99461	0.99477	0.99492	0.99506	0.9952
2.6	0.99534	0.99547	0.9956	0.99573	0.99585	0.99598	0.99609	0.99621	0.99632	0.99643
2.7	0.99653	0.99664	0.99674	0.99683	0.99693	0.99702	0.99711	0.9972	0.99728	0.99736
2.8	0.99744	0.99752	0.9976	0.99767	0.99774	0.99781	0.99788	0.99795	0.99801	0.99807
2.9	0.99813	0.99819	0.99825	0.99831	0.99836	0.99841	0.99846	0.99851	0.99856	0.99861
3.0	0.99865	0.99869	0.99874	0.99878	0.99882	0.99886	0.99889	0.99893	0.99896	0.999

Solve for n

Roadmap: Bounding Error

Goal: Find the value of n such that 98% of the time, the estimate \bar{X} is within 5% of the true p

1. Define probability of a “bad event”
2. Apply CLT
3. Convert to a standard normal
4. Solve for n

Idealized Polling

So far, we have been discussing “idealized polling”. Real life is normally not so nice 😞

Assumed we can sample people uniformly at random, not really possible in practice

- Not everyone responds
- Response rates might differ in different groups
- Will people respond truthfully?

Makes polling in real life much more complex than this idealized model!