

No activity slide today

# Expectation

CSE 312 Spring 21  
Lecture 11

# Announcements

*Feedback*  
HW2 was just released

There's a post on Ed about a common bug in HW2 P6. Please read that before filing a regrade request on that problem.

HW3 due tonight

Remember you have late days. Particularly if you run into debugging issues with the programming part.

HW4 out (late) tonight

*Kushal will return on Friday*

# Try it yourself

"pmf"  $f_X(x) = \mathbb{P}(X=x)$

What is the CDF of  $X$  where

$X$  be the largest value among the three balls. (Drawing 3 of the 20 without replacement)

$\rightarrow F_X(x) = \mathbb{P}(X \leq x)$

$\mathbb{P}(X \leq x) = \mathbb{P}(\max \text{ of three} \leq x) = \mathbb{P}(\text{each of the three is } \leq x)$

$\hat{f}$   
 $x \in 4, 3 \leq x \leq 20$

$$= \frac{\binom{x}{3}}{\binom{20}{3}}$$

# Try it yourself

What is the CDF of  $X$  where

$X$  be the largest value among the three balls. (Drawing 3 of the 20 without replacement)

$$\mathbb{P}(X \leq -3) = 0$$

$$\mathbb{P}(X \leq 25)$$

$$F_X(x) = \begin{cases} 0 & \text{if } x < 3 \\ \binom{[x]}{3} / \binom{20}{3} & \text{if } 3 \leq x \leq 20 \\ 1 & \text{otherwise} \end{cases}$$

$$\mathbb{P}(X \leq 5.5) = \mathbb{P}(X \leq 5)$$

# Try it yourself

What is the CDF of  $X$  where

$X$  be the largest value among the three balls. (Drawing 3 of the 20 without replacement)

$$\underbrace{P(X \leq k)} \leq \underbrace{P(X \leq k+3)}$$

$$F_X(x) = \begin{cases} 0 & \text{if } x < 3 \\ \binom{\lfloor x \rfloor}{3} / \binom{20}{3} & \text{if } 3 \leq x \leq 20 \\ 1 & \text{otherwise} \end{cases}$$

otherwise

Good checks: Is  $F_X(\infty) = 1$ ? If not, something is wrong.

Is  $F_X(x)$  increasing? If not something is wrong.

Is  $F_X(x)$  defined for all real number inputs? If not something is wrong.

# Two descriptions

## PROBABILITY MASS FUNCTION

Defined for all  $\mathbb{R}$  inputs.

Usually has "0 otherwise" as an extra case.

$$\sum_x f_X(x) = 1$$

$$0 \leq f_X(x) \leq 1$$

$$\sum_{z:z \leq x} f_X(z) = F_X(x)$$

## CUMULATIVE DISTRIBUTION FUNCTION

Defined for all  $\mathbb{R}$  inputs.


Usually has "0 otherwise" and 1 otherwise" extra cases

Non-decreasing function

$$0 \leq F_X(x) \leq 1$$

$$\lim_{x \rightarrow -\infty} F_X(x) = 0$$

$$\lim_{x \rightarrow \infty} F_X(x) = 1$$



**Expectation**

---



# Expectation

$$\mathbb{E}[X]$$

## Expectation

The “expectation” (or “expected value”) of a random variable  $X$  is:

$$\mathbb{E}[X] = \sum_k k \cdot \mathbb{P}(X = k)$$

Intuition: The weighted average of values  $X$  could take on.

Weighted by the probability you actually see them.



# Example 1

Flip a fair coin twice (independently)

Let  $X$  be the number of heads.

$$\mathbb{P}[X=x] = f(x)$$

$$\left. \begin{array}{l} \\ \\ \\ \\ \end{array} \right\} \begin{array}{l} 1/4 \\ 1/2 \\ 1/4 \\ 0 \end{array}$$

$$\text{if } x=0$$

$$\text{if } x=1$$

$$\text{if } x=2$$

$$\text{o/w}$$

$\Omega = \{\underline{TT}, \underline{TH}, \underline{HT}, \underline{HH}\}$ ,  $\mathbb{P}()$  is uniform measure.

$$\underline{\mathbb{E}[X]} = \frac{1}{4} \cdot 0 + \frac{1}{2} \cdot 1 + \frac{1}{4} \cdot 2 = 0 + \frac{1}{2} + \frac{1}{2} = \underline{1}.$$

# Example 2

You roll a biased die.

It shows a 6 with probability  $\frac{1}{3}$ , and 1, ..., 5 with probability  $\frac{2}{15}$  each.

Let  $X$  be the value of the die. What is  $\mathbb{E}[X]$ ?

$$\begin{aligned} & \frac{1}{3} \cdot 6 + \frac{2}{15} \cdot 5 + \frac{2}{15} \cdot 4 + \frac{2}{15} \cdot 3 + \frac{2}{15} \cdot 2 + \frac{2}{15} \cdot 1 \\ & = 2 + \frac{2(5+4+3+2+1)}{15} = 2 + \frac{30}{15} = 4 \end{aligned}$$

$\mathbb{E}[X]$  is not just the most likely outcome!

# Try it yourself

Let  $X$  be the result shown on a fair die. What is  $\mathbb{E}[X]$ ?

Let  $Y$  be the sum of two (independent) fair die rolls. What is  $\mathbb{E}[Y]$ ?

Fill out the poll everywhere so Robbie  
knows how long to explain  
Go to [pollev.com/cse312](https://pollev.com/cse312)

# Try it yourself

Let  $X$  be the result shown on a fair die. What is  $\mathbb{E}[X]$

$$\begin{aligned} & 6 \cdot \frac{1}{6} + 5 \cdot \frac{1}{6} + 4 \cdot \frac{1}{6} + 3 \cdot \frac{1}{6} + 2 \cdot \frac{1}{6} + 1 \cdot \frac{1}{6} \\ &= \frac{21}{6} = \underline{3.5} \end{aligned}$$

$\mathbb{E}[X]$  is not necessarily a possible outcome!

That's ok, it's an average!

# Try it yourself

$$\mathbb{E}[Y] =$$

$$\frac{1}{36} \cdot 2 + \frac{2}{36} \cdot 3 + \frac{3}{36} \cdot 4 + \frac{4}{36} \cdot 5 + \frac{5}{36} \cdot 6 + \frac{6}{36} \cdot 7 + \frac{5}{36} \cdot 8 + \frac{4}{36} \cdot 9 + \frac{3}{36} \cdot 10 + \frac{2}{36} \cdot 11 + \frac{1}{36} \cdot 12$$

$$= 7$$

$\mathbb{E}[Y] = 2\underbrace{\mathbb{E}[X]}$ . That's not a coincidence...we'll talk about why next time.

# Subtle but Important

$X$  is random. You don't know what it is (at least until you run the experiment).

~~$\mathbb{E}[X]$~~  is not random. It's a number.

You don't need to run the experiment to know what it is.



**More Independence**

---

# Independence of events

Recall the definition of independence of **events**:

## Independence

Two events  $A, B$  are independent if  
 $\mathbb{P}(A \cap B) = \mathbb{P}(A) \cdot \mathbb{P}(B)$

'if  $\mathbb{P}(A) > 0$ ,  $\mathbb{P}(B) > 0$ ,  
then  $\Leftrightarrow \mathbb{P}(A|B) = \mathbb{P}(A) \Leftrightarrow \mathbb{P}(B|A) = \mathbb{P}(B)$ .



# Independence for 3 or more events

$$\mathbb{P}(A_1 \cap \dots \cap A_n) = \mathbb{P}(A_1) \mathbb{P}(A_2 | A_1) \mathbb{P}(A_3 | A_1 \cap A_2) \dots \mathbb{P}(A_n | \dots)$$

For three or more events, we need two kinds of independence

## Pairwise Independence

Events  $A_1, A_2, \dots, A_n$  are pairwise independent if

$$\mathbb{P}(A_i \cap A_j) = \mathbb{P}(A_i) \cdot \mathbb{P}(A_j) \text{ for all } i, j \quad i \neq j.$$

## Mutual Independence

Events  $A_1, A_2, \dots, A_n$  are mutually independent if

$$\mathbb{P}(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = \mathbb{P}(A_{i_1}) \cdot \mathbb{P}(A_{i_2}) \cdots \mathbb{P}(A_{i_k})$$

for every subset  $\{i_1, i_2, \dots, i_k\}$  of  $\{1, 2, \dots, n\}$ .


# Pairwise Independence vs. Mutual Independence

Roll two fair dice (one red one blue) independently

$R$  = "red die is 3"

$B$  = "blue die is 5"

$S$  = "sum is 7"



How should we describe these events?

# Pairwise Independence

$R, B, S$  are pairwise independent

$$\mathbb{P}(R \cap B) \stackrel{?}{=} \mathbb{P}(R)\mathbb{P}(B)$$

$$\frac{1}{36} = \frac{1}{6} \cdot \frac{1}{6} \text{ Yes! (These are also independent by the problem statement)}$$

$$\mathbb{P}(R \cap S) \stackrel{?}{=} \mathbb{P}(R)\mathbb{P}(S)$$

$$\frac{1}{36} \stackrel{?}{=} \frac{1}{6} \cdot \frac{1}{6} \text{ Yes!}$$

$$\mathbb{P}(B \cap S) \stackrel{?}{=} \mathbb{P}(B)\mathbb{P}(S)$$

$$\frac{1}{36} \stackrel{?}{=} \frac{1}{6} \cdot \frac{1}{6} \text{ Yes!}$$

Since all three pairs are independent, we say the random variables are pairwise independent.

# Mutual Independence

$R, B, S$  are not mutually independent.

→  $\mathbb{P}(R \cap B \cap S) = 0$ ; if the red die is 3, and blue die is 5 then the sum is 8 (so it can't be 7)

$$\mathbb{P}(R)\mathbb{P}(B)\mathbb{P}(S) = \left(\frac{1}{6}\right)^3 = \frac{1}{216} \neq 0$$

# Checking Mutual Independence

It's not enough to check just  $\mathbb{P}(A \cap B \cap C)$  either.

Roll a fair 8-sided die.

Let  $A$  be  $\{1,2,3,4\}$

$B$  be  $\{2,4,6,8\}$

$C$  be  $\{2,3,5,7\}$

$$\mathbb{P}(A \cap B \cap C) = \mathbb{P}(\{2\}) = \frac{1}{8}$$

$$\mathbb{P}(A)\mathbb{P}(B)\mathbb{P}(C) = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}$$

# Checking Mutual Independence

It's not enough to check just  $\mathbb{P}(A \cap B \cap C)$  either.

Roll a fair 8-sided die.

Let  $A$  be  $\{1,2,3,4\}$

$B$  be  $\{2,4,6,8\}$

$C$  be  $\{2,3,5,7\}$

$$\mathbb{P}(A \cap B \cap C) = \mathbb{P}(\{2\}) = \frac{1}{8}$$

$$\mathbb{P}(A)\mathbb{P}(B)\mathbb{P}(C) = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8}$$

But  $A$  and  $B$  aren't independent (nor are  $B, C$ ; though  $A$  and  $C$  are independent). Because there's a subset that's not independent,  $A, B, C$  are not mutually independent.

# Checking Mutual Independence

To check mutual independence of events:

Check **every** subset.

To check pairwise independence of events:

Check **every** subset of size two.

# Independence of Random Variables

That's for events...what about random variables?

## Independence (of random variables)

$X$  and  $Y$  are independent if for all  $k, \ell$

$$\mathbb{P}(X = k, Y = \ell) = \mathbb{P}(X = k)\mathbb{P}(Y = \ell)$$

We'll often use commas instead of  $\cap$  symbol.



# Independence of Random Variables

The “for all values” is important.

We say that the event “the sum is 7” is independent of “the red die is 5”  
What about  $S$  = “the sum of two dice” and  $R$  = “the value of the red die”

# Independence of Random Variables

The “for all values” is important.

We say that the event “the sum is 7” is independent of “the red die is 5”  
What about  $S$  = “the sum of two dice” and  $R$  = “the value of the red die”

NOT independent.

$\mathbb{P}(S = 2, R = 5) \neq \mathbb{P}(S = 2)\mathbb{P}(R = 5)$  (for example)

# Independence of Random Variables

Flip a coin independently  $2n$  times.

Let  $X$  be "the number of heads in the first  $n$  flips."

Let  $Y$  be "the number of heads in the last  $n$  flips."

$X$  and  $Y$  are independent.

# Mutual Independence for RVs

A little simpler to write down than for events


## Mutual Independence (of random variables)

$X_1, X_2, \dots, X_n$  are mutually independent if for all  $x_1, x_2, \dots, x_n$

$$\mathbb{P}(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) = \mathbb{P}(X_1 = x_1)\mathbb{P}(X_2 = x_2) \cdots \mathbb{P}(X_n = x_n)$$

DON'T need to check all subsets for random variables...

But you do need to check all values (all possible  $x_i$ ) still.



## Extra Practice

---

# More Practice

Suppose you flip a coin until you see a heads for the first time.

Let  $X$  be the number of trials (including the heads)

What is the pmf of  $X$ ?

The cdf of  $X$ ?

$\mathbb{E}[X]$ ?

# More Practice

Suppose you flip a coin until you see a heads for the first time.

Let  $X$  be the number of trials (including the heads)

What is the pmf of  $X$ ?  $f_X(x) = 1/2^x$  for  $x \in \mathbb{Z}^+$ , 0 otherwise

The cdf of  $X$ ?  $F_X(x) = 1 - 1/2^{\lfloor x \rfloor}$  for  $x \geq 0$ , 0 for  $x < 0$ .

$$\mathbb{E}[X]? \sum_{i=1}^{\infty} \frac{i}{2^i} = 2$$

# More Random Variable Practice

Roll a fair die  $n$  times. Let  $X$  be the number of rolls that are 5s or 6s.

What is the pmf?

Don't try to write the CDF...it's a mess...

Or try for a few minutes to realize it isn't nice.

What is the expectation?



# More Random Variable Practice

Roll a fair die  $n$  times. Let  $Z$  be the number of rolls that are 5s or 6s.

What's the probability of getting exactly  $k$  5's/6's? Well we need to know which  $k$  of the  $n$  rolls are 5's/6's. And then multiply by the probability of getting exactly that outcome

$$f_Z(z) = \begin{cases} \binom{n}{z} \cdot \left(\frac{1}{3}\right)^z \left(\frac{2}{3}\right)^{n-z} & \text{if } z \in Z, 0 \leq z \leq n \\ 0 & \text{otherwise} \end{cases}$$

Expectation formula is a mess. If you plug it into a calculator you'll get a nice, clean simplification:  $n/3$ .