

**CSE 312**

# **Foundations of Computing II**

**Lecture 5: Conditional Probability and Bayes Theorem**

## Review Probability

**Definition.** A **sample space**  $\Omega$  is the set of all possible outcomes of an experiment.

**Definition.** An **event**  $E \subseteq \Omega$  is a subset of possible outcomes.

*~*

Examples:

- Single coin flip:  $\Omega = \{H, T\}$
- Two coin flips:  $\Omega = \{HH, HT, TH, TT\}$
- Roll of a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$

Examples:

- Getting at least one head in two coin flips:  
 $E = \{HH, HT, TH\}$
- Rolling an even number on a die :  
 $E = \{2, 4, 6\}$

## Review Probability space

Either finite or infinite countable (e.g., integers)

**Definition.** A (discrete) **probability space** is a pair  $(\Omega, P)$  where:

- $\Omega$  is a set called the **sample space**.
- $P$  is the **probability measure**, a function  $P: \Omega \rightarrow \mathbb{R}$  such that:

- $P(x) \geq 0$  for all  $x \in \Omega$
- $\sum_{x \in \Omega} P(x) = 1$

Set of possible elementary outcomes

$$A \subseteq \Omega: P(A) = \sum_{x \in A} P(x)$$

Specify Likelihood (or probability) of each elementary outcome

Some outcome must show up


The likelihood (or probability) of each outcome is non-negative.

# Agenda

- Conditional Probability ◀
- Bayes Theorem
- Law of Total Probability
- More Examples



## Conditional Probability (Idea)



$$P(\text{ice cream}) = \frac{43}{70}$$

What's the probability that someone likes ice cream **given** they like donuts?

$\text{ice cream \& donuts}$

$\text{donuts}$

$$\frac{7}{7 + 13} = \frac{7}{20}$$

## Conditional Probability

**Definition.** The **conditional probability** of event  $A$  **given** an event  $B$  happened (assuming  $P(B) \neq 0$ ) is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

"probability of  $A$  given  $B$ "

An equivalent and useful formula is

$$P(A \cap B) = P(A|B)P(B)$$

## Conditional Probability Examples

Suppose that you flip a fair coin twice.


What is the probability that both flips are heads given that you have at least one head?

Let  $O$  be the event that at least one flip is heads

Let  $B$  be the event that both flips are heads

$$P(O) = \frac{3}{4} \quad P(B) = \frac{1}{4} \quad P(B \cap O) = \frac{1}{4}$$

$$P(B|O) = \frac{P(B \cap O)}{P(O)} = \frac{\frac{1}{4}}{\frac{3}{4}} = \frac{1}{3}$$



## Conditional Probability Examples


Suppose that you flip a fair coin twice.

What is the probability that at least one flip is heads given that at least one flip is tails?

$$P(H|T) = \frac{P(H \cap T)}{P(T)} = \frac{1/4}{3/4} = \frac{1}{3}$$

Let  $H$  be the event that at least one flip is heads

Let  $T$  be the event that at least one flip is tails





## Conditional Probability Examples

Suppose that you flip a fair coin twice.

What is the probability that at least one flip is heads given that at least one flip is tails?

Let  $H$  be the event that at least one flip is *heads*

Let  $T$  be the event that at least one flip is *tails*

$$\underline{P(H) = 3/4} \quad P(T) = 3/4 \quad P(H \cap T) = 1/2$$

$$P(H|T) = \frac{P(H \cap T)}{P(T)} = \frac{1/2}{3/4} = \frac{2}{3}$$

$\Omega$

HH	HT
TH	TT

## Reversing Conditional Probability

**Question:** Does  $P(A|B) = P(B|A)$ ?

No!

- Let  $A$  be the event you are wet
- Let  $B$  be the event you are swimming


$$P(A|B) = 1$$
$$P(B|A) \neq 1$$

## Example with Conditional Probability

pollev.com/paulbeame028

Suppose we toss a red die and a blue die:  
both 6 sided and all outcomes equally  
likely.

What is  $P(B)$ ? What is  $P(B|A)$ ?



	$P(B)$	$P(B A)$	
a)	1/6	1/6	$\times P(A B)$
$\rightarrow$ b)	1/6	1/3	$\checkmark$
c)	1/6	<del>3/36</del>	<del><math>P(A)</math></del>
d)	1/9	1/3	$\checkmark$

$\uparrow$   
 $\frac{6}{36}$   
 $\frac{1}{3}$

## Gambler's fallacy

Assume we toss 51 fair coins.

Assume we have seen 50 coins, and they are all "tails".

What are the odds the 51<sup>st</sup> coin is "heads"?

$A$  = first 50 coins are "tails"

$$\frac{1}{2^{50}}$$

$B$  = first 50 coins are "tails", 51<sup>st</sup> coin is "heads"

$$\frac{1}{2^{51}}$$

$$Pr[B|A] = Pr(B)$$
$$= Pr(A \cap B)$$

51<sup>st</sup> coin is independent of (A)  
outcomes of first 50 tosses!


$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{1/2^{51}}{1/2^{50}} = \frac{1}{2}$$

**Gambler's fallacy** = Feels like it's time for "heads"!?

# Agenda

- Conditional Probability
- Bayes Theorem ◀
- Law of Total Probability
- More Examples

# Bayes Theorem



A formula to let us “reverse” the conditional.

**Theorem. (Bayes Rule)** For events  $A$  and  $B$ , where  $P(A), P(B) > 0$ ,

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

$P(A)$  is called the **prior** (our belief without knowing anything)

$P(A|B)$  is called the **posterior** (our belief after learning  $B$ )

## Bayes Theorem Proof

Claim:

$$P(A), P(B) > 0 \Rightarrow P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

$$P(A \cap B) = P(A|B)P(B)$$

$$P(B \cap A) = P(B|A)P(A)$$

$$\therefore P(A|B)P(B) = P(B|A)P(A)$$

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

since  
 $P(B) \neq 0$   
divide by  $P(B)$

## Bayes Theorem Proof

Claim:

$$P(A), P(B) > 0 \Rightarrow P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

By definition of conditional probability

$$P(A \cap B) = P(A|B)P(B)$$

Swapping  $A, B$  gives

$$P(B \cap A) = P(B|A)P(A)$$

But  $P(A \cap B) = P(B \cap A)$ , so


$$P(A|B)P(B) = P(B|A)P(A)$$

Dividing both sides by  $P(B)$  gives

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$



# Brain Break




# Agenda

- Conditional Probability
- Bayes Theorem
- Law of Total Probability ◀
- More Examples

## Partitions (Idea)

These events **partition** the sample space

1. They “cover” the whole space
2. They don’t overlap





## Partition

**Definition.** Non-empty events  $E_1, E_2, \dots, E_n$  **partition** the sample space  $\Omega$  if  
(Exhaustive)

$$E_1 \cup E_2 \cup \dots \cup E_n = \bigcup_{i=1}^n E_i = \Omega$$

(Pairwise Mutually Exclusive)


$$\forall_i \forall_{i \neq j} E_i \cap E_j = \emptyset$$



## Law of Total Probability (Idea)

If we know  $E_1, E_2, \dots, E_n$  partition  $\Omega$ , what can we say about  $P(F)$ ?

*What is  $P(F)$ ?*



## Law of Total Probability (LTP)

**Definition.** If events  $E_1, E_2, \dots, E_n$  partition the sample space  $\Omega$ , then for any event  $F$

$$\underline{P(F)} = \underline{P(F \cap E_1)} + \dots + \underline{P(F \cap E_n)} = \sum_{i=1}^n \underline{P(F \cap E_i)}$$

$$P(F|E_i)P(E_i)$$

Using the definition of conditional probability  $P(F \cap E) = P(F|E)P(E)$

We can get the alternate form of this that shows

$$\underline{P(F)} = \sum_{i=1}^n \underbrace{P(F|E_i)}_{\text{easy}} \underbrace{P(E_i)}_{\text{easy}}$$

## Another Contrived Example


Alice has two pockets:

- **Left pocket:** Two blue balls, two green balls
- **Right pocket:** One blue ball, two green balls.

Alice picks a random ball from a random pocket.

[Both pockets equally likely, each ball equally likely.]

## Sequential Process



- **Left pocket:** Two blue, two green
- **Right pocket:** One blue, two green

$$1/3 = P(\text{B}|\text{Right}) \text{ and } 2/3 = P(\text{G}|\text{Right})$$

$$P(\text{B}) = P(\text{B} \cap \text{Left}) + P(\text{B} \cap \text{Right}) \quad (\text{Law of total probability})$$

$$= P(\text{Left}) \times P(\text{B}|\text{Left}) + P(\text{Right}) \times P(\text{B}|\text{Right})$$

$$= \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{3} = \frac{1}{4} + \frac{1}{6} = \frac{5}{12}$$



## Agenda

- Conditional Probability
- Bayes Theorem
- Law of Total Probability
- **More Examples** ◀

## Example – Zika Testing

Zika fever

OVERVIEW SYMPTOMS SPECIALISTS

Fever  
Rash  
Joint pain  
Red eyes



Spread through mosquito bites

Source

A disease caused by Zika virus that's spread through mosquito bites.

The image shows a medical information card for Zika fever. At the top, it says 'Zika fever' and has three tabs: 'OVERVIEW', 'SYMPTOMS', and 'SPECIALISTS'. Under the 'SYMPTOMS' tab, a list of symptoms is shown: 'Fever', 'Rash', 'Joint pain', and 'Red eyes'. Below the list is an illustration of a woman with a red rash on her chest and a mosquito biting her arm. A circular inset shows a close-up of a mosquito on skin. Below the illustration, it says 'Spread through mosquito bites' and 'Source'. At the bottom, a text box states: 'A disease caused by Zika virus that's spread through mosquito bites.'

Usually no or mild symptoms (rash); sometimes severe symptoms (paralysis).

During pregnancy: may cause birth defects.

Suppose you took a Zika test, and it returns “positive”, what is the likelihood that you actually have the disease?

- Tests for diseases are rarely 100% accurate.

## Example – Zika Testing

Suppose we know the following Zika stats

- A test is 98% effective at detecting Zika (“true positive”)  $P(T|Z)$
- However, the test may yield a “false positive” 1% of the time  $P(T|Z^c)$
- 0.5% of the US population has Zika.  $P(Z)$

What is the probability you have Zika (event  $Z$ ) if you test positive (event  $T$ )?.