

CSE 312

Foundations of Computing II

Lecture 3: Even more counting

Binomial Theorem, Inclusion-Exclusion, Pigeonhole Principle

Recap

Two core rules for counting a set S :

- **Sum rule:**
 - Break up S into **disjoint** pieces/cases
 - $|S|$ = the **sum** of the sizes of the pieces.
- **Product rule:**
 - View the elements of S as being constructed by a **series of choices**, where the # of possibilities for each choice doesn't depend on the previous choices
 - $|S|$ = the **product** of the # of choices in each step of the series.

Recap

- **k -sequences**: How many length k sequences over alphabet of size n ?
 - Product rule $\rightarrow n^k$
- **k -permutations**: How many length k sequences over alphabet of size n , **without repetition**?
 - Permutation $\rightarrow \frac{n!}{(n-k)!}$
- **k -combinations**: How many size k subsets of a set of size n (**without repetition and without order**)?
 - Combination $\rightarrow \binom{n}{k} = \frac{n!}{k!(n-k)!}$

Binomial Coefficients – Many interesting and useful properties

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$$\binom{n}{n} = 1$$

$$\binom{n}{1} = n$$

$$\binom{n}{0} = 1$$

Fact. $\binom{n}{k} = \binom{n}{n-k}$

Symmetry in Binomial Coefficients

Fact. $\sum_{k=0}^n \binom{n}{k} = 2^n$

Follows from Binomial Theorem

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

Pascal's Identity

Agenda

- Binomial Theorem
- Combinatorial Proofs for Pascal Identity ◀
- Inclusion-Exclusion
- Pigeonhole Principle
- Counting Practice

Pascal's Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

How to prove Pascal's identity?

Algebraic argument:

$$\begin{aligned}\binom{n-1}{k-1} + \binom{n-1}{k} &= \frac{(n-1)!}{(k-1)!(n-k)!} + \frac{(n-1)!}{k!(n-1-k)!} \\ &= 20 \text{ years later ...} \\ &= \frac{n!}{k!(n-k)!} \\ &= \binom{n}{k}\end{aligned}$$


Hard work and not intuitive

Let's see a combinatorial argument

Example – Pascal's Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S| = |B| + |A|$



$S = A \cup B$

Combinatorial proof idea:

- Find *disjoint* sets A and B such that A , B , and $S = A \cup B$ have the sizes above.
- The equation then follows by the Sum Rule.

One Minute Discussion

Example – Pascal's Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$$|S| = |B| + |A|$$

Combinatorial proof idea:

- Find *disjoint* sets A and B such that A , B , and $S = A \cup B$ have these sizes

$$|S| = \binom{n}{k}$$

S : set of size k subsets of $[n] = \{1, 2, \dots, n\}$.

e.g. $n = 4, k = 2$, $S = \{\{1,2\}, \{1,3\}, \{1,4\}, \{2,3\}, \{2,4\}, \{3,4\}\}$

A : set of size k subsets of $[n - 1]$ (i.e., DON'T include n)

$$A = \{\{1,2\}, \{1,3\}, \{2,3\}\}$$

B : Choose a size $k - 1$ subsets of $[n - 1]$ then add n (i.e., DO include n)

$$B = \{\{1,4\}, \{2,4\}, \{3,4\}\}$$

Example – Pascal's Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S| = |B| + |A|$

S : set of size k subsets of $[n] = \{1, 2, \dots, n\}$.

A : set of size k subsets of $[n-1]$ (i.e., DON'T include n)

B : Choose a size $k-1$ subsets of $[n-1]$ then add n

Combinatorial proof idea:

- Find *disjoint* sets A and B such that A, B , and $S = A \cup B$ have these sizes

n not in set, need to choose k elements from $[n-1]$

$$|B| = \binom{n-1}{k}$$

n is in set, need to choose other $k-1$ elements from $[n-1]$


$$|A| = \binom{n-1}{k-1}$$

Agenda

- Binomial Theorem
- Combinatorial Proofs
- Inclusion-Exclusion ◀
- Pigeonhole Principle
- Counting Practice

Recap Disjoint Sets


Sets that do not contain common elements ($A \cap B = \emptyset$)



$$\text{Sum Rule: } |A \cup B| = |A| + |B|$$

Inclusion-Exclusion

But what if the sets are not disjoint?




$$\begin{aligned} |A| &= 43 \\ |B| &= 20 \\ |A \cap B| &= 7 \\ |A \cup B| &= ??? \end{aligned}$$

Fact. $|A \cup B| = |A| + |B| - |A \cap B|$

Inclusion-Exclusion

Not drawn to scale

What if there are three sets?



$$\begin{aligned} |A| &= 43 \\ |B| &= 20 \\ |C| &= 35 \\ |A \cap B| &= 7 \\ |A \cap C| &= 16 \\ |B \cap C| &= 11 \\ |A \cap B \cap C| &= 4 \\ |A \cup B \cup C| &= ??? \end{aligned}$$

Fact.

$$\begin{aligned} |A \cup B \cup C| &= |A| + |B| + |C| \\ &\quad - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C| \end{aligned}$$

Inclusion-Exclusion

Let A, B be sets. Then

$$|A \cup B| = |A| + |B| - |A \cap B|$$

In general, if A_1, A_2, \dots, A_n are sets, then

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_n| &= \textit{singles} - \textit{doubles} + \textit{triples} - \textit{quads} + \dots \\ &= (|A_1| + \dots + |A_n|) - (|A_1 \cap A_2| + \dots + |A_{n-1} \cap A_n|) + \dots \end{aligned}$$

Agenda


- Binomial Theorem
- Combinatorial Proofs
- Inclusion-Exclusion
- Pigeonhole Principle ◀
- Counting Practice

Pigeonhole Principle (PHP): Idea

10 pigeons, 9 pigeonholes



Pigeonhole Principle: Idea



If 11 children have to share 3 cakes, at least one cake must be shared by how many children?

Pigeonhole Principle – More generally

If there are n pigeons in $k < n$ holes, then one hole must contain at least $\frac{n}{k}$ pigeons!

Proof. Assume there are $< \frac{n}{k}$ pigeons per hole.

Then, there are $< k \cdot \frac{n}{k} = n$ pigeons overall.

Contradiction!

Pigeonhole Principle – Better version

If there are n pigeons in $k < n$ holes, then one hole must contain at least $\left\lceil \frac{n}{k} \right\rceil$ pigeons!

Reason. Can't have fractional number of pigeons

Syntax reminder:

- Ceiling: $\lceil x \rceil$ is x rounded up to the nearest integer (e.g., $\lceil 2.731 \rceil = 3$)
- Floor: $\lfloor x \rfloor$ is x rounded down to the nearest integer (e.g., $\lfloor 2.731 \rfloor = 2$)

Pigeonhole Principle: Strategy

To use the PHP to solve a problem, there are generally 4 steps

1. Identify pigeons
2. Identify pigeonholes
3. Specify how pigeons are assigned to pigeonholes
4. Apply PHP

Pigeonhole Principle – Example

In a room with 367 people, there are at least two with the same birthday.

Solution:

1. **367** pigeons = people
2. **366** holes (365 for a normal year + Feb 29) = possible birthdays
3. Person goes into hole corresponding to own birthday
4. By PHP, there must be two people with the same birthday

Pigeonhole Principle – Example (Surprising?)

*In every set S of 100 integers, there are at least **two** elements whose difference is a multiple of 37.*

When solving a PHP problem:

1. Identify pigeons
2. Identify pigeonholes
3. Specify how pigeons are assigned to pigeonholes
4. Apply PHP

One Minute Discussion

Pigeonhole Principle – Example (Surprising?)

*In every set S of 100 integers, there are at least **two** elements whose difference is a multiple of 37.*

When solving a PHP problem:

1. Identify pigeons
2. Identify pigeonholes
3. Specify how pigeons are assigned to pigeonholes
4. Apply PHP

Pigeons: integers x in S

Pigeonholes: $\{0,1,\dots,36\}$


Assignment: x goes to $x \bmod 37$

Since $100 > 37$, by PHP, there are $x \neq y \in S$ s.t.
 $x \bmod 37 = y \bmod 37$ which implies
 $x - y = 37k$ for some integer k

Agenda

- Binomial Theorem
- Combinatorial Proofs
- Inclusion-Exclusion
- Pigeonhole Principle
- Counting Practice ◀

Quick Review of Cards



How many possible 5 card hands?


$$\binom{52}{5}$$

- 52 total cards
- 13 different ranks: 2,3,4,5,6,7,8,9,10,J,Q,K,A
- 4 different suits: Hearts, Diamonds, Clubs, Spades

Counting Cards I

- 52 total cards
- 13 different ranks: 2,3,4,5,6,7,8,9,10,J,Q,K,A
- 4 different suits: Hearts, Diamonds, Clubs, Spades

- A "straight" is five consecutive rank cards of any suit (where A,2,3,4,5 also counts as consecutive).
How many possible straights?



$$10 \cdot 4^5 = 10,240$$

Counting Cards II

- 52 total cards
- 13 different ranks: 2,3,4,5,6,7,8,9,10,J,Q,K,A
- 4 different suits: Hearts, Diamonds, Clubs, Spades

- A flush is five card hand all of the same suit.
How many possible flushes?



$$4 \cdot \binom{13}{5} = 5148$$

Counting Cards III

- 52 total cards
- 13 different ranks: 2,3,4,5,6,7,8,9,10,J,Q,K,A
- 4 different suits: Hearts, Diamonds, Clubs, Spades

- A flush is five card hand all of the same suit.
How many possible flushes?


$$4 \cdot \binom{13}{5} = 5148$$



- How many flushes are **NOT** straights?

= #flush - #flush and straight

$$\left(4 \cdot \binom{13}{5} = 5148 \right) - 10 \cdot 4$$



Sleuth's Criterion (Rudich)

For each object constructed, it should be possible to reconstruct the **unique** sequence of choices that led to it.


No sequence → under counting Many sequences → over counting

EXAMPLE: How many ways are there to choose a 5 card hand that contains at least 3 Aces?

First choose 3 Aces. Then choose remaining two cards.

$$\binom{4}{3} \cdot \binom{49}{2}$$

Poll:

- A. Correct
- B. Overcount 
- C. Undercount

[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)

<https://pollev.com/paulbeame028>

Sleuth's Criterion (Rudich)

For each object constructed, it should be possible to reconstruct the **unique** sequence of choices that led to it.

Many sequences → over counting

EXAMPLE: How many ways are there to choose a hand that contains at least 3 Aces?

First choose 3 Aces. Then choose remaining two cards.

$$\binom{4}{3} \cdot \binom{49}{2}$$

Problem: This counts a hand with all 4 Aces in 4 different ways!

e.g. it counts $A\clubsuit, A\spadesuit, A\heartsuit, A\diamonds, 2\heartsuit$

four times:

$\{A\clubsuit, A\spadesuit, A\heartsuit\} \{A\diamonds, 2\heartsuit\}$

$\{A\clubsuit, A\spadesuit, A\diamonds\} \{A\heartsuit, 2\heartsuit\}$

$\{A\clubsuit, A\heartsuit, A\spadesuit\} \{A\diamonds, 2\heartsuit\}$

$\{A\spadesuit, A\heartsuit, A\diamonds\} \{A\clubsuit, 2\heartsuit\}$

Sleuth's Criterion (Rudich)

For each object constructed, it should be possible to reconstruct the **unique** sequence of choices that led to it.

No sequence → under counting Many sequences → over counting

EXAMPLE: How many ways are there to choose a 5 card hand that contains at least 3 Aces?

Use the sum rule

= # 5 card hand containing exactly 3 Aces

+ # 5 card hand containing exactly 4 Aces

$$\binom{4}{3} \cdot \binom{48}{2}$$

$$\binom{48}{1}$$

Counting when order only *partly* matters

We often want to count # of partly ordered lists:

Let M = # of ways to produce fully ordered lists

P = # of partly ordered lists

N = # of ways to produce corresponding fully ordered list given a partly ordered list


Then $M = P \cdot N$ by the product rule. Often M and N are easy to compute:

$$P = M/N$$

Dividing by N “removes” part of the order.

Rooks on chessboard

How many ways to place two identical rooks on a chessboard so that they don't share a row or a column



Fully ordered: Pretend Rooks are different

1. Column for rook1
2. Row for rook1
3. Column for rook2
4. Row for rook2

$$(8 \cdot 7)^2$$

“Remove” the order of the two rooks:

$$(8 \cdot 7)^2 / 2$$

Binomial Theorem

Theorem. Let $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$ a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

Set $x = y = 1$

Corollary.

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

Binomial Theorem: A less obvious consequence

Theorem. Let $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$ a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

$= -1$ if k is odd
 $= +1$ if k is even

Corollary. For every n , if O and E are the sets of odd and even integers between 0 and n

$$\sum_{k \in O} \binom{n}{k} = \sum_{k \in E} \binom{n}{k}$$

e.g., $n=4$: 1 4 6 4 1

Proof: Set $x = -1, y = 1$ in the binomial theorem

Tools and concepts

- Sum rule, Product rule
- Permutations, combinations
- Inclusion-exclusion
- Binomial Theorem
- Combinatorial proofs
- Pigeonhole principle
- Binary encoding/stars and bars