

CSE 312

Foundations of Computing II

Lecture 2: Permutations, combinations, the Binomial Theorem and more.




Anna R. Karlin

Slide Style Credit: Based on Stefano Tessaro's slides for 312 19au incorporating ideas from Alex Tsun, Rachel Lin, Hunter Schafer & myself 😊

Grading, syllabus and administrivia

- Questions?

Agenda

- Recap & Examples 
- Binomial Theorem
- Multinomial Coefficients
- Inclusion-Exclusion
- Combinatorial Proofs

<https://pollev.com/annakarlin185>

Quick Summary

- **Sum Rule**

If you can choose from

- **Either** one of n options,
- **OR** one of m options with **NO overlap** with the previous n ,

then the number of possible outcomes of the experiment is $n + m$

- **Product Rule**

In a sequential process, if there are

- n_1 choices for the first step,
- n_2 choices for the second step (given the first choice), ..., and
- n_k choices for the k^{th} step (given the previous choices),

then the total number of outcomes is $n_1 \times n_2 \times \cdots \times n_k$





- **Complementary Counting**

Quick Summary

- **K-sequences**: How many length k sequences over alphabet of size n ?
repetition allowed.
 - Product rule $\rightarrow n^k$
- **K-permutations**: How many length k sequences over alphabet of size n , without repetition?
 - Permutation $\rightarrow \frac{n!}{(n-k)!}$
- **K-combinations**: How many size k subsets of a set of n distinct elements (without repetition and without order)?
 - Combination $\rightarrow \binom{n}{k} = \frac{n!}{k!(n-k)!}$

Product rule – Another example

5 books



“How many ways are there to distribute 5 books among Alice, Bob, and Charlie?”

Every book to one person, everyone gets ≥ 0 books.



Alice




Bob



Charlie

Example Book Assignment



Book assignment – Modeling

Correct?


Poll:

- A. right
- B. Overcount
- C. Undercount
- D. No idea

<https://pollev.com/annakarlin185>


$2^5 = 32$ options

λ



$A = \{\text{orange book}, \text{blue book}\}$


$2^5 = 32$ options



$B = \{\text{green book}\}$

$2^5 = 32$ options

λ





$C = \{\text{blue book}, \text{grey book}\}$


= 32^3 assignment

Problem – Overcounting

Problem: We are counting some invalid assignments!!!
→ overcounting!



What went wrong in the sequential process?
After assigning set A to Alice, set B is no longer a valid option for Bob



Book assignment – Second try

$2^5 = 32$ options

χ



$A = \{\text{orange book}, \text{blue book}\}$

χ







$B = \{\text{green book}\}$



$C = \{\text{blue book}, \text{brown book}\}$

Product rule – A better way

5 books



“How many ways are there to distribute 5 books among Alice, Bob, and Charlie?”

Every book to one person, everyone gets ≥ 0 books.



Alice



Bob



Charlie






Book assignments – Choices tell you who gets each book

X

X

X

X




Lesson: Representation of what we are counting is very important!

Think about the various possible ways you could make a sequence of choices that leads to an outcome in the set of outcomes you are trying to count.




Example – Counting Paths



“How many ways to walk from 1st and Spring to 5th and Pine only going \uparrow and \rightarrow ?”

Example – Counting Paths -2



“How many ways to walk from 1st and Spring to 5th and Pine only going \uparrow and \rightarrow ?”

Poll:

A. 2^7

B. $\frac{7!}{4!}$

C. $\binom{7}{4} = \frac{7!}{4!3!}$

D. $\binom{7}{3} = \frac{7!}{3!4!}$

<https://pollev.com/annakarlin185>

Symmetry in Binomial Coefficients

Fact. $\binom{n}{k} = \binom{n}{n-k}$

Proof. $\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n!}{(n-k)!k!} = \binom{n}{n-k}$

Why??




This is called an Algebraic proof, i.e., Prove by checking algebra

Symmetry in Binomial Coefficients – A different proof


Fact. $\binom{n}{k} = \binom{n}{n-k}$

Two **equivalent** ways to choose k out of n objects (unordered)

1. Choose which k elements are **included**
2. Choose which $n - k$ elements are **excluded**



$$\binom{4}{1} = 4 = \binom{4}{3}$$



Symmetry in Binomial Coefficients – A different proof

Fact. $\binom{n}{k} = \binom{n}{n-k}$

Two **equivalent** ways to choose k out of n objects (unordered)


1. Choose which k elements are **included**
2. Choose which $n - k$ elements are **excluded**

This is called a **combinatorial argument/proof**

- Let S be a set of objects
- Show how to count $|S|$ one way $\Rightarrow |S| = N$
- Show how to count $|S|$ another way $\Rightarrow |S| = m$


More examples of
combinatorial proofs
coming soon!

Example – Counting Paths - 3



“How many ways to walk from 1st and Spring to 5th and Pine only going \uparrow and \rightarrow but stopping at Starbucks on 3rd and Pike?”

Example – Counting Paths - 3



“How many ways to walk from 1st and Spring to 5th and Pine only going \uparrow and \rightarrow but stopping at Starbucks on 3rd and Pike?”

Poll:

A. $\binom{7}{3}$

B. $\binom{7}{3} \binom{7}{1}$

C. $\binom{4}{2} \binom{3}{1}$

D. $\binom{4}{2} \binom{3}{2}$

<https://pollev.com/annakarlin185>

Agenda

- Recap & Examples
- **Binomial Theorem** ◀
- Multinomial Coefficients
- Inclusion-Exclusion
- Combinatorial Proofs

Binomial Theorem: Idea

$$\begin{aligned}(x + y)^2 &= (x + y)(x + y) \\ &= xx + xy + yx + yy \\ &= x^2 + 2xy + y^2\end{aligned}$$

$$\begin{aligned}(x + y)^4 &= (x + y)(x + y)(x + y)(x + y) \\ &= xxxx + yyyy + xyxy + yxyy + \dots\end{aligned}$$

Binomial Theorem: Idea

Poll: What is the coefficient for xy^3 ?

- A. 4
- B. $\binom{4}{1}$
- C. $\binom{4}{3}$
- D. 3

<https://pollev.com/annakarlin185>

$$(x + y)^4 = (x + y)(x + y)(x + y)(x + y)$$
$$= xxxx + yyyy + xyxy + yxyy + \dots$$

Binomial Theorem: Idea

$$(x + y)^n = (x + y)(x + y)(x + y) \cdots (x + y)$$

Each term is of the form $x^k y^{n-k}$, since each term is made by multiplying exactly n variables, either x or y .

How many times do we get $x^k y^{n-k}$?

Binomial Theorem: Idea

$$(x + y)^n = (x + y)(x + y)(x + y) \cdots (x + y)$$

Each term is of the form $x^k y^{n-k}$, since each term is made by multiplying exactly n variables, either x or y .

How many times do we get $x^k y^{n-k}$? The number of ways to choose k of the n variables we multiply to be an x (the rest will be y).

$$\binom{n}{k} = \binom{n}{n-k}$$

Binomial Theorem

Theorem. Let $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$ a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

Binomial Theorem

Theorem. Let $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$ a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

Corollary.

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

Agenda

- Recap & Examples
- Binomial Theorem
- **Multinomial Coefficients** ◀
- Inclusion-Exclusion
- Combinatorial Proofs

Example – Word Permutations

How many ways to re-arrange the letters in the word “MATH”?

Poll:

A. $\binom{26}{4}$

B. 4^4

C. $4!$

D. I don't know



MATH

<https://pollev.com/annakarlin185>

Example – Word Permutations

How many ways to re-arrange the letters in the word “MUUMUU”?



Example – Word Permutations

How many ways to re-arrange the letters in the word “MUUMUU”?



Choose where the 2 M's go, and then the U's are set **OR**
Choose where the 4 U's go, and then the M's are set

Either way, we get $\binom{6}{2} \cdot \binom{4}{4} = \binom{6}{4} \cdot \binom{2}{2} = \frac{6!}{2!4!}$

Another way to think about it

How many ways to re-arrange the letters in the word “MUUMUU”?

Arrange the 6 letters as if they were distinct.

$$M_1 U_1 U_2 M_2 U_3 U_4$$

Then divide by $4!$ to account for duplicate M's and divide by $2!$ to account for duplicate U's.

Yields $\frac{6!}{2!4!}$



Another example – Word Permutations

How many ways to re-arrange the letters in the word “GODOGGY”?



Poll:

A. $7!$

B. $\frac{7!}{3!}$

C. $\frac{7!}{3!2!1!1!}$

D. $\binom{7}{3} \cdot \binom{5}{2} \cdot 3!$

--	--	--	--	--	--	--

<https://pollev.com/annakarlin185>

Multinomial coefficients

If we have k types of objects, with n_1 of the first type, n_2 of the second type, ..., n_k of the k^{th} type, where $n = n_1 + n_2 + \cdots + n_k$ then the number of arrangements of the n objects is

$$\binom{n}{n_1, n_2, \dots, n_k} = \frac{n!}{n_1! n_2! \cdots n_k!}$$

Note that objects of the same type are indistinguishable.

Example – Word Permutations

How many ways to re-arrange the letters in the word “GODOGGY”?




$n = 7$ (length of sequence) $K = 4$ types = $\{G, O, D, Y\}$

$n_1 = 3, n_2 = 2, n_3 = 1, n_4 = 1$


$$\binom{6}{4,2,1,1} = \frac{6!}{2!4!1!1!}$$

Agenda

- Recap & Examples
- Binomial Theorem
- Multinomial Coefficients
- **Inclusion-Exclusion** 
- Combinatorial Proofs

Recap Disjoint Sets


Sets that do not contain common elements ($A \cap B = \emptyset$)



Sum Rule: $|A \cup B| = |A| + |B|$

Inclusion-Exclusion

But what if the sets are not disjoint?



$$|A| = 43$$

$$|B| = 20$$

$$|A \cap B| = 7$$


$$|A \cup B| = ???$$

Fact. $|A \cup B| = |A| + |B| - |A \cap B|$

Inclusion-Exclusion

Not drawn to scale

What if there are three sets?



$$|A| = 43$$

$$|B| = 20$$

$$|C| = 35$$

$$|A \cap B| = 7$$

$$|A \cap C| = 16$$

$$|B \cap C| = 11$$

$$|A \cap B \cap C| = 4$$

$$|A \cup B \cup C| = ???$$

Fact.

$$\begin{aligned} |A \cup B \cup C| &= |A| + |B| + |C| \\ &\quad - |A \cap B| - |A \cap C| - |B \cap C| \\ &\quad + |A \cap B \cap C| \end{aligned}$$

Inclusion-Exclusion

Let A, B be sets. Then

$$|A \cup B| = |A| + |B| - |A \cap B|$$

In general, if A_1, A_2, \dots, A_n are sets, then

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_n| &= \textit{singles} - \textit{doubles} + \textit{triples} - \textit{quads} + \dots \\ &= (|A_1| + \dots + |A_n|) - (|A_1 \cap A_2| + \dots + |A_{n-1} \cap A_n|) + \dots \end{aligned}$$

Agenda

- Recap & Examples
- Binomial Theorem
- Multinomial Coefficients
- Inclusion-Exclusion
- **Combinatorial Proofs** ◀

Combinatorial proof: Show that $M = N$

- Let S be a set of objects
- Show how to count $|S|$ one way $\Rightarrow |S| = M$
- Show how to count $|S|$ another way $\Rightarrow |S| = N$
- Conclude that $M = N$

Binomial Coefficient – Many interesting and useful properties

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$$\binom{n}{n} = 1$$

$$\binom{n}{1} = n$$

$$\binom{n}{0} = 1$$

Fact. $\binom{n}{k} = \binom{n}{n-k}$

Symmetry in Binomial Coefficients

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

Pascal's Identity

Fact. $\sum_{k=0}^n \binom{n}{k} = 2^n$

Follows from Binomial theorem

Pascal's Identities

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

How to prove Pascal's identity?

Algebraic argument:

$$\begin{aligned}\binom{n-1}{k-1} + \binom{n-1}{k} &= \frac{(n-1)!}{(k-1)!(n-k)!} + \frac{(n-1)!}{k!(n-1-k)!} \\ &= 20 \text{ years later ...} \\ &= \frac{n!}{k!(n-k)!} \\ &= \binom{n}{k}\end{aligned}$$


Hard work and not intuitive

Let's see a combinatorial argument

Example – Binomial Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S| = |A| + |B|$



$S = A \cup B$, disjoint

S : the set of size k subsets of $[n] = \{1, 2, \dots, n\} \rightarrow |S| = \binom{n}{k}$

A : the set of size k subsets of $[n]$ including n


B : the set of size k subsets of $[n]$ NOT including n

Sum rule:
 $|A \cup B| = |A| + |B|$

Example – Binomial Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S| = |A| + |B|$



S: the set of size k subsets of $[n] = \{1, 2, \dots, n\}$ $\rightarrow |S| = \binom{n}{k}$

e.g.: $n = 4$, $S = \{\{1,2\}, \{1,3\}, \{1,4\}, \{2,3\}, \{2,4\}, \{3,4\}\}$

A: the set of size k subsets of $[n]$ including n

$$A = \{\{1,4\}, \{2,4\}, \{3,4\}\}. \quad n = 4$$

B: the set of size k subsets of $[n]$ NOT including n

$$B = \{\{1,2\}, \{1,3\}, \{2,3\}\}$$


Example – Binomial Identity

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

\uparrow \uparrow \uparrow

$|S|$ $|A|$ $|B|$

$S = A \cup B$



S : the set of size k subsets of $[n] = \{1, 2, \dots, n\}$

A : the set of size k subsets of $[n]$ including n

B : the set of size k subsets of $[n]$ NOT including n

n is in set, need to choose $k - 1$ elements from $[n - 1]$

$$|A| = \binom{n-1}{k-1}$$

n not in set, need to choose k elements from $[n - 1]$

$$|B| = \binom{n-1}{k}$$

combinatorial argument/proof

- Elegant
- Simple
- Intuitive



[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)

Algebraic argument

- Brute force
- Less Intuitive



[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)