

CSE 312

# Foundations of Computing II

Lecture 4: Intro to discrete probability



**Anna R. Karlin**


Style Credit: Based on Stefano Tessaro's slides for 312 19au  
incorporating ideas from Alex Tsun, Rachel Lin, Hunter Schafer & myself ☺

Plus few slides from Berkeley CS 70

# Probability

- We want to model uncertainty.
  - i.e., outcome not determined a-priori
  - E.g. throwing dice, flipping a coin...
  - We want to numerically measure likelihood of outcomes = probability.
  - We want to make complex statements about these likelihoods.
- We will not argue why a certain physical process realizes the probabilistic model we study
  - Why is the outcome of the coin flip really “random”?
- First part of class: “Discrete” probability theory
  - Experiment with finite / discrete set of outcomes.
  - Will explore countably infinite and continuous outcomes later

## Agenda

- Events 
- Probability
- Equally Likely Outcomes
- Probability Axioms and Beyond Equally Likely Outcomes
- More Examples

## Sample Space

**Definition.** A **sample space**  $\Omega$  is the set of all possible outcomes of an experiment.

omega

### Examples:

- Single coin flip:  $\Omega = \{\underline{H}, \underline{T}\}$
- Two coin flips:  $\Omega = \{\underline{HH}, \underline{HT}, \underline{TH}, \underline{TT}\}$
- Roll of a die:  $\Omega = \{1, 2, 3, 4, 5, 6\}$

## Events


**Definition.** An event  $E \subseteq \Omega$  is a subset of possible outcomes.

↑  
sample space

### Examples:

- Getting at least one head in two coin flips:  $E = \{HH, HT, TH\}$
- Rolling an even number on a die :  $E = \{2, 4, 6\}$

## Events



**Definition.** An **event**  $E \subseteq \Omega$  is a subset of possible outcomes.

### Examples:

- Getting at least one head in two coin flips:  $E = \{HH, HT, TH\}$
- Rolling an even number on a die :  $E = \{2, 4, 6\}$

**Definition.** Events  $E$  and  $F$  are **mutually exclusive** if  $E \cap F = \emptyset$  (i.e., can't happen at same time)

### Examples:

- For dice rolls: If  $E = \{2, 4, 6\}$  and  $F = \{1, 5\}$ , then  $E \cap F = \emptyset$

## Example: 4-sided Dice

$$|\Omega| = 16$$

Suppose I roll two 4-sided dice Let  $D_1$  be the value of the blue die and  $D_2$  be the value of the red die. To the right is the sample space (possible outcomes).

What outcomes match these events?

A.  $D_1 = 1$

B.  $D_1 + D_2 = 6$

C.  $D_1 = 2 * D_2$

Die 1 ( $D_1$ )

Die 2 ( $D_2$ )

	1	2	3	4
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)

## Example: 4-sided Dice

Suppose I roll two 4-sided dice Let  $D_1$  be the value of the blue die and  $D_2$  be the value of the red die. To the right is the sample space (possible outcomes).

What outcomes match these events?

A.  $D_1 = 1$

$$A = \{(1,1), (1,2), (1,3), (1,4)\}$$

B.  $D_1 + D_2 = 6$

$$B = \{(2,4), (3,3), (4,2)\}$$

C.  $D_1 = 2 * D_2$

$$C = \{(2,1), (4,2)\}$$

Die 1 ( $D_1$ )

Die 2 ( $D_2$ )

	1	2	3	4
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)



## Example: 4-sided Dice, Mutual Exclusivity

Are  $A$  and  $B$  mutually exclusive?  
How about  $B$  and  $C$ ?

A.  $D_1 = 1$

B.  $D_1 + D_2 = 6$

C.  $D_1 = 2 * D_2$

Die 1 ( $D_1$ )

Die 2 ( $D_2$ )

	1	2	3	4
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)

# Agenda

- Events
- **Probability** ◀
- Equally Likely Outcomes
- Probability Axioms and Beyond Equally Likely Outcomes
- More Examples

## Idea: Probability

A **probability** is a number (between 0 and 1) describing how likely a particular outcome will happen.

Will define a function

$$\mathbb{P}: \Omega \rightarrow [0, 1]$$

that maps outcomes  $\omega \in \Omega$  to probabilities.

– Also use notation:  $\mathbb{P}(\omega)$  =  $P(\omega)$  =  $\text{Pr}(\omega)$

## Example – Coin Tossing


Imagine we toss one coin – outcome can be **heads** or **tails**.

$$\underline{\Omega = \{H, T\}}$$

$\mathbb{P}$ ? Depends! What do we want to model?!

Fair coin toss

$$\underline{\mathbb{P}(H) = \mathbb{P}(T) = \frac{1}{2} = 0.5}$$



## Example – Coin Tossing


Imagine we toss one coin – outcome can be **heads** or **tails**.

$$\Omega = \{H, T\}$$

$\mathbb{P}$ ? Depends! What do we want to model?!

**Bent** coin toss (e.g., biased or unfair coin)

$$\mathbb{P}(H) = 0.45, \quad \mathbb{P}(T) = 0.55$$




## Probability space

**Definition.** A (discrete) **probability space** is a pair  $(\Omega, \mathbb{P})$  where:

- $\Omega$  is a set called the sample space.
- $\mathbb{P}$  is the **probability measure**,

a function  $\mathbb{P}: \Omega \rightarrow [0,1]$  such that:

- $\mathbb{P}(\omega) \geq 0$  for all  $\omega \in \Omega$
- $\sum_{\omega \in \Omega} \mathbb{P}(\omega) = 1$



# Probability space

Either finite or infinite countable (e.g., integers)


**Definition.** A (discrete) **probability space** is a pair  $(\Omega, \mathbb{P})$  where:

- $\Omega$  is a set called the **sample space**.
- $\mathbb{P}$  is the **probability measure**, a function  $\mathbb{P}: \Omega \rightarrow [0,1]$  such that:

- $\mathbb{P}(\omega) \geq 0$  for all  $\omega \in \Omega$

- $\sum_{\omega \in \Omega} \mathbb{P}(\omega) = 1$

Set of possible elementary outcomes



Specify Likelihood (or probability) of each elementary outcome

Some outcome must show up

The likelihood (or probability) of each outcome is non-negative.

## Uniform Probability Space

**Definition.** A uniform probability space is a pair  $(\Omega, \mathbb{P})$  such that

$$\mathbb{P}(x) = \frac{1}{|\Omega|}$$

for all  $x \in \Omega$ .

Examples:

- Fair coin  $P(x) = \frac{1}{2}$
- Fair 6-sided die  $P(x) = \frac{1}{6}$




$$P(\omega) = \frac{1}{|\Omega|}$$

## Events

**Definition.** An **event** in a probability space  $(\Omega, \mathbb{P})$  is a subset  $\mathcal{A} \subseteq \Omega$ . Its probability is

$$\underline{\mathbb{P}(\mathcal{A})} = \sum_{\omega \in \mathcal{A}} \mathbb{P}(\omega)$$



Convenient abuse of notation:  $\mathbb{P}$  is extended to be defined over **sets**.

$$\mathbb{P}(\omega) = \mathbb{P}(\{\omega\})$$



## Agenda

- Events
- Probability
- **Equally Likely Outcomes** ◀
- Probability Axioms and Beyond Equally Likely Outcomes
- More Examples

$$|\Omega| = 16$$

$$\Pr(\omega) = \frac{1}{16}$$

## Example: 4-sided Dice, Event Probability

Think back to 4-sided die. Suppose each die is fair. What is the probability of event  $B$ ?  $\Pr(B) = ???$

$$B. D1 + D2 = 6$$

$$B = \{(2,4), (3,3), (4,2)\}$$

$$\begin{aligned} \Pr(B) &= \Pr((2,4)) + \Pr((3,3)) \\ &\quad + \Pr((4,2)) \\ &= \frac{3}{16} \end{aligned}$$

Die 1 (D1)

Die 2 (D2)

	1	2	3	4
1	(1, 1)	(1, 2)	(1, 3)	(1, 4)
2	(2, 1)	(2, 2)	(2, 3)	(2, 4)
3	(3, 1)	(3, 2)	(3, 3)	(3, 4)
4	(4, 1)	(4, 2)	(4, 3)	(4, 4)

uniform P.S.

$$\Pr(E) = \sum_{\omega \in E} \Pr(\omega)$$

$$P(\omega) = \frac{1}{|\Omega|} \quad \forall \omega \in \Omega$$

## Equally Likely Outcomes

If  $(\Omega, P)$  is a **uniform** probability space, then for any event  $E \subseteq \Omega$ , then

$$P(E) = \frac{|E|}{|\Omega|}$$

This follows from the definitions of the prob. of an event and uniform probability spaces.

$$Pr(E) = \sum_{\omega \in E} Pr(\omega) = \sum_{\omega \in E} \left( \frac{1}{|\Omega|} \right) = \frac{|E|}{|\Omega|}$$

$$\Omega = \{\text{sequences of } 100 \text{ coin flips}\} = \underline{\underline{\{H, T\}^{100}}}$$

3

### Example – Coin Tossing

HHT, HHT, HTH, HTT, ...

$$|\Omega| = 2^{100}$$

Toss a coin 100 times. Each outcome is equally likely. What is the probability of seeing 50 heads?

uniform prob space.


$$\Pr(\omega) = \frac{1}{2^{100}}$$

<https://pollev.com/annakarlin185>

- (A)  $\frac{1}{2}$
- (B)  $\frac{1}{2^{50}}$
- (C)  $\frac{\binom{100}{50}}{2^{100}}$
- (D) Not sure

$$\Pr(\text{seeing exactly 50 heads}) = \frac{|\{\text{outcomes with 50 heads}\}|}{2^{100}}$$

$$= \frac{\binom{100}{50}}{2^{100}}$$



## Brain Break



## Agenda

- Events
- Probability
- Equally Likely Outcomes
- **Probability Axioms and Beyond Equally Likely Outcomes** ◀
- More Examples

$$|\Omega| = n$$

# possible events  $2^n$

## Axioms of Probability


Let  $\Omega$  denote the sample space and  $E, F \subseteq \Omega$  be events. Note this applies to **any** probability space (not just uniform)

$$P: \Omega \rightarrow [0, 1]$$

**Axiom 1 (Non-negativity):**  $P(E) \geq 0$ .

**Axiom 2 (Normalization):**  $P(\Omega) = 1$

**Axiom 3 (Countable Additivity):** If  $E$  and  $F$  are mutually exclusive, then  $P(E \cup F) = P(E) + P(F)$



**Corollary 1 (Complementation):**  $P(E^c) = 1 - P(E)$ .


**Corollary 2 (Monotonicity):** If  $E \subseteq F$ ,  $P(E) \leq P(F)$

**Corollary 3 (Inclusion-Exclusion):**  $P(E \cup F) = P(E) + P(F) - P(E \cap F)$

$$P(E^c) = 1 - P(E)$$

Cor 2

$$\begin{aligned} \Omega &= E \cup E^c \\ \uparrow & \quad \uparrow \\ 1 &= P(E) + P(E^c) \end{aligned}$$




0.333

$$\begin{aligned} E^c &= \overline{E} \\ &= \Omega \setminus E \end{aligned}$$



car 3




## Review Probability space

Either finite or infinite countable (e.g., integers)

**Definition.** A (discrete) **probability space** is a pair  $(\Omega, \mathbb{P})$  where:

- $\Omega$  is a set called the **sample space**.
- $\mathbb{P}$  is the **probability measure**, a function  $\mathbb{P}: \Omega \rightarrow [0,1]$  such that:
  - $\mathbb{P}(\omega) \geq 0$  for all  $\omega \in \Omega$
  - $\sum_{\omega \in \Omega} \mathbb{P}(\omega) = 1$

Set of possible elementary outcomes





Specify Likelihood (or probability) of each elementary outcome

Some outcome must show up

The likelihood (or probability) of each outcome is non-negative.



## Non-equally Likely Outcomes

Probability spaces can have **non-equally likely outcomes**.



$\{HH, HT, TH, TT\}$   
0    $\frac{1}{2}$     $\frac{1}{2}$    0

# More Examples of Non-equally Likely Outcomes



## Agenda

- Events
- Probability
- Equally Likely Outcomes
- Probability Axioms and Beyond Equally Likely Outcomes
- **More Examples** ◀

## Example: Dice Rolls

red die      green die

Suppose I had a two, fair, 6-sided dice that we roll once each. What is the probability that we see *at least one 3* in the two rolls.

$$\Omega = \{ \text{all pairs of die rolls} \}$$

$$|\Omega| = 36$$

$$\Pr(\omega) = \frac{1}{36}$$

$$E = \{ \text{at least one 3} \}$$

$$\Pr(E) = \frac{|E|}{|\Omega|} = \frac{|\Omega| - |\text{no 3's}|}{36} = \frac{36 - 5^2}{36}$$

$$= 1 - \Pr(E^c)$$

$$= 1 - \frac{25}{36}$$

no 3's

$$\frac{|E^c|}{|\Omega|} = \frac{25}{36}$$

## Example: Birthday "Paradox"

Suppose we have a collection of  $n$  people in a room. What is the probability that at least 2 people share a birthday? Assuming there are 365 possible birthdays, with uniform probability for each day.

$\{A, B, C, D, \dots\}$   
 $\frac{1}{365}$   $\frac{317}{365}$   $\frac{4}{365}$

$\Omega = \{ \text{set of birthdays of } n \text{ people} \}$

$$|\Omega| = 365^n$$

$$\Pr(\omega) = \frac{1}{365^n}$$

$$\Pr(\text{at least 2 people share a bday}) = 1 - \Pr(\text{no 2 people share a bday})$$

$$= 1 - \frac{365 \cdot 364 \cdot 363 \cdots (365 - n + 1)}{365^n} \ll$$

Example: Birthday "Paradox" cont.

Out of  $n$  people  
 Pr (at least one of  $n$  people has same birthday as me)  
 Out of

$n = 23$   
 $Pr(E) > 0.5$   
 $n = 100$   
 $Pr(E) > 0.98$

$$= 1 - Pr(\text{none of } n \text{ people have oct 17})$$



$n = 23$	0.06	$\frac{\text{\# of ways w/ no Oct 17}}{365^n} = \frac{364^n}{365^n}$
$n = 100$	0.23	