

**CSE 312**

# **Foundations of Computing II**

**Lecture 2:** Permutations, combinations, the Binomial Theorem and more.


## Grading, syllabus and administrivia

- Please be sure that you read the syllabus carefully!!!!
- Questions?

## Survival Tips

- Don't fall behind.
  - Do every single concept check.
  - Do the reading (better yet, ahead of time!)
  - Go through the section problems. Try to solve them. Make sure you understand the solutions.
- Take the homework seriously.
  - Get started early.
  - Come to office hours if you need help.
  - Form study groups!
  - Take academic integrity seriously.

# Agenda

- Recap & Examples 
- Binomial Theorem
- Multinomial Coefficients
- Inclusion-Exclusion
- Combinatorial Proofs

<https://pollev.com/annakarlin185>

## Quick Summary

- **Sum Rule**

If you can choose from

- **Either** one of  $n$  options,
- **OR** one of  $m$  options with **NO overlap** with the previous  $n$ ,

then the number of possible outcomes of the experiment is  $n + m$

- **Product Rule**

In a sequential process, if there are

- $n_1$  choices for the first step,
- $n_2$  choices for the second step (given the first choice), ..., and
- $n_k$  choices for the  $k^{\text{th}}$  step (given the previous choices),

then the total number of outcomes is  $n_1 \times n_2 \times \cdots \times n_k$





- **Complementary Counting**

## Quick Summary

- **K-sequences**: How many length  $k$  sequences over alphabet of size  $n$ ?  
repetition allowed.
  - Product rule  $\rightarrow n^k$
- **K-permutations**: How many length  $k$  sequences over alphabet of size  $n$ , without repetition?
  - Permutation  $\rightarrow \frac{n!}{(n-k)!}$
- **K-combinations**: How many size  $k$  subsets of a set of  $n$  distinct elements (without repetition and without order)?
  - Combination  $\rightarrow \binom{n}{k} = \frac{n!}{k!(n-k)!}$

## Product rule – Another example

5 books



*“How many ways are there to distribute 5 books among Alice, Bob, and Charlie?”*

Every book to one person, everyone gets  $\geq 0$  books.



Alice




Bob



Charlie

# Example Book Assignment





## Book assignment – Modeling

**Correct?**


Poll:

- A. right
- B. Overcount
- C. Undercount
- D. No idea

<https://pollev.com/annakarlin185>

$2^5 = 32$  options

λ




$A = \{\text{orange book}, \text{blue book}\}$

$2^5 = 32$  options



$B = \{\text{green book}\}$

$2^5 = 32$  options





$C = \{\text{blue book}, \text{grey book}\}$


=  $32^3$  assignment

## Problem – Overcounting

**Problem:** We are counting some invalid assignments!!!  
→ overcounting!




What went wrong in the sequential process?  
After assigning set  $A$  to Alice, set  $B$  is no longer a valid option for Bob



## Book assignment – Second try

$2^5 = 32$  options

$\chi$



$A = \{\text{orange book}, \text{blue book}\}$

$\chi$







$B = \{\text{green book}\}$



$C = \{\text{blue book}, \text{grey book}\}$

## Product rule – A better way

5 books



*“How many ways are there to distribute 5 books among Alice, Bob, and Charlie?”*

Every book to one person, everyone gets  $\geq 0$  books.



Alice



Bob



Charlie





# Book assignments – Choices tell you who gets each book

X

X

X

X




# ***Lesson: Representation of what we are counting is very important!***

**Think about the various possible ways you could make a sequence of choices that leads to an outcome in the set of outcomes you are trying to count.**




## Example – Counting Paths



*“How many ways to walk from 1<sup>st</sup> and Spring to 5<sup>th</sup> and Pine only going  $\uparrow$  and  $\rightarrow$  ?”*

## Example – Counting Paths -2



*“How many ways to walk from 1<sup>st</sup> and Spring to 5<sup>th</sup> and Pine only going  $\uparrow$  and  $\rightarrow$  ?”*

Poll:

A.  $2^7$

B.  $\frac{7!}{4!}$

C.  $\binom{7}{4} = \frac{7!}{4!3!}$

D.  $\binom{7}{3} = \frac{7!}{3!4!}$

<https://pollev.com/annakarlin185>



## Symmetry in Binomial Coefficients

**Fact.**  $\binom{n}{k} = \binom{n}{n-k}$

## Symmetry in Binomial Coefficients

**Fact.**  $\binom{n}{k} = \binom{n}{n-k}$

**Proof.**  $\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n!}{(n-k)!k!} = \binom{n}{n-k}$

**Why??**




This is called an Algebraic proof, i.e., Prove by checking algebra

## Symmetry in Binomial Coefficients – A different proof


**Fact.**  $\binom{n}{k} = \binom{n}{n-k}$

Two **equivalent** ways to choose  $k$  out of  $n$  objects (unordered)

1. Choose which  $k$  elements are **included**
2. Choose which  $n - k$  elements are **excluded**



$$\binom{4}{1} = 4 = \binom{4}{3}$$



## Symmetry in Binomial Coefficients – A different proof

**Fact.**  $\binom{n}{k} = \binom{n}{n-k}$

Two **equivalent** ways to choose  $k$  out of  $n$  objects (unordered)


1. Choose which  $k$  elements are **included**
2. Choose which  $n - k$  elements are **excluded**

This is called a **combinatorial argument/proof**

- Let  $S$  be a set of objects
- Show how to count  $|S|$  one way  $\Rightarrow |S| = N$
- Show how to count  $|S|$  another way  $\Rightarrow |S| = m$


More examples of  
combinatorial proofs  
coming soon!

## Example – Counting Paths - 3



*“How many ways to walk from 1<sup>st</sup> and Spring to 5<sup>th</sup> and Pine only going  $\uparrow$  and  $\rightarrow$  but stopping at Starbucks on 3<sup>rd</sup> and Pike?”*

## Example – Counting Paths - 3



*“How many ways to walk from 1<sup>st</sup> and Spring to 5<sup>th</sup> and Pine only going  $\uparrow$  and  $\rightarrow$  but stopping at Starbucks on 3<sup>rd</sup> and Pike?”*

Poll:

A.  $\binom{7}{3}$

B.  $\binom{7}{3} \binom{7}{1}$

C.  $\binom{4}{2} \binom{3}{1}$

D.  $\binom{4}{2} \binom{3}{2}$

<https://pollev.com/annakarlin185>

## Agenda

- Recap & Examples
- **Binomial Theorem** ◀
- Multinomial Coefficients
- Inclusion-Exclusion
- Combinatorial Proofs

## Binomial Theorem: Idea

$$\begin{aligned}(x + y)^2 &= (x + y)(x + y) \\ &= xx + xy + yx + yy \\ &= x^2 + 2xy + y^2\end{aligned}$$

$$\begin{aligned}(x + y)^4 &= (x + y)(x + y)(x + y)(x + y) \\ &= xxxx + yyyy + xyxy + yxyy + \dots\end{aligned}$$



## Binomial Theorem: Idea

Poll: What is the coefficient for  $xy^3$ ?

- A. 4
- B.  $\binom{4}{1}$
- C.  $\binom{4}{3}$
- D. 3

<https://pollev.com/annakarlin185>

$$(x + y)^4 = (x + y)(x + y)(x + y)(x + y)$$
$$= xxxx + yyyy + xyxy + yxyy + \dots$$

## Binomial Theorem: Idea

$$(x + y)^n = (x + y)(x + y)(x + y) \cdots (x + y)$$

Each term is of the form  $x^k y^{n-k}$ , since each term is made by multiplying exactly  $n$  variables, either  $x$  or  $y$ .

How many times do we get  $x^k y^{n-k}$ ?

## Binomial Theorem: Idea

$$(x + y)^n = (x + y)(x + y)(x + y) \cdots (x + y)$$

Each term is of the form  $x^k y^{n-k}$ , since each term is made by multiplying exactly  $n$  variables, either  $x$  or  $y$ .

How many times do we get  $x^k y^{n-k}$ ? The number of ways to choose  $k$  of the  $n$  variables we multiply to be an  $x$  (the rest will be  $y$ ).

$$\binom{n}{k} = \binom{n}{n-k}$$

## Binomial Theorem

**Theorem.** Let  $x, y \in \mathbb{R}$  and  $n \in \mathbb{N}$  a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

## Binomial Theorem

**Theorem.** Let  $x, y \in \mathbb{R}$  and  $n \in \mathbb{N}$  a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

**Corollary.**

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

## Agenda

- Recap & Examples
- Binomial Theorem
- **Multinomial Coefficients** ◀
- Inclusion-Exclusion
- Combinatorial Proofs

## Example – Word Permutations

*How many ways to re-arrange the letters in the word “MATH”?*

Poll:

A.  $\binom{26}{4}$

B.  $4^4$

C.  $4!$

D. I don't know



MATH

<https://pollev.com/annakarlin185>

## Example – Word Permutations

*How many ways to re-arrange the letters in the word “MUUMUU”?*





## Example – Word Permutations

*How many ways to re-arrange the letters in the word “MUUMUU”?*



Choose where the 2 M's go, and then the U's are set **OR**  
Choose where the 4 U's go, and then the M's are set

Either way, we get  $\binom{6}{2} \cdot \binom{4}{4} = \binom{6}{4} \cdot \binom{2}{2} = \frac{6!}{2!4!}$

## Another way to think about it

*How many ways to re-arrange the letters in the word “MUUMUU”?*

Arrange the 6 letters as if they were distinct.

$$M_1 U_1 U_2 M_2 U_3 U_4$$

Then divide by  $4!$  to account for duplicate M's and divide by  $2!$  to account for duplicate U's.

Yields  $\frac{6!}{2!4!}$



## Another example – Word Permutations

*How many ways to re-arrange the letters in the word “GODOGGY”?*



Poll:

A.  $7!$

B.  $\frac{7!}{3!}$

C.  $\frac{7!}{3!2!1!1!}$

D.  $\binom{7}{3} \cdot \binom{4}{2} \cdot 2!$

--	--	--	--	--	--	--

<https://pollev.com/annakarlin185>

## Multinomial coefficients

If we have  $k$  types of objects, with  $n_1$  of the first type,  $n_2$  of the second type, ...,  $n_k$  of the  $k^{\text{th}}$  type, where  $n = n_1 + n_2 + \cdots + n_k$  then the number of arrangements of the  $n$  objects is

$$\binom{n}{n_1, n_2, \dots, n_k} = \frac{n!}{n_1! n_2! \cdots n_k!}$$

Note that objects of the same type are indistinguishable.

## Example – Word Permutations

*How many ways to re-arrange the letters in the word “GODOGGY”?*




$n = 7$  (length of sequence)     $K = 4$  types =  $\{G, O, D, Y\}$

$n_1 = 3, n_2 = 2, n_3 = 1, n_4 = 1$


$$\binom{6}{4,2,1,1} = \frac{6!}{2!4!1!1!}$$

## Agenda

- Recap & Examples
- Binomial Theorem
- Multinomial Coefficients
- **Inclusion-Exclusion** 
- Combinatorial Proofs

## Recap Disjoint Sets


Sets that do not contain common elements ( $A \cap B = \emptyset$ )



**Sum Rule:**  $|A \cup B| = |A| + |B|$

## Inclusion-Exclusion

But what if the sets are not disjoint?



$$|A| = 43$$

$$|B| = 20$$

$$|A \cap B| = 7$$

$$|A \cup B| = ???$$


**Fact.**  $|A \cup B| = |A| + |B| - |A \cap B|$



# Inclusion-Exclusion

Not drawn to scale

What if there are three sets?



$$|A| = 43$$

$$|B| = 20$$

$$|C| = 35$$

$$|A \cap B| = 7$$

$$|A \cap C| = 16$$

$$|B \cap C| = 11$$

$$|A \cap B \cap C| = 4$$

$$|A \cup B \cup C| = ???$$

**Fact.**

$$\begin{aligned} |A \cup B \cup C| &= |A| + |B| + |C| \\ &\quad - |A \cap B| - |A \cap C| - |B \cap C| \\ &\quad + |A \cap B \cap C| \end{aligned}$$

## Inclusion-Exclusion

Let  $A, B$  be sets. Then

$$|A \cup B| = |A| + |B| - |A \cap B|$$

In general, if  $A_1, A_2, \dots, A_n$  are sets, then

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_n| &= \text{singles} - \text{doubles} + \text{triples} - \text{quads} + \dots \\ &= (|A_1| + \dots + |A_n|) - (|A_1 \cap A_2| + \dots + |A_{n-1} \cap A_n|) + \dots \end{aligned}$$

## Agenda

- Recap & Examples
- Binomial Theorem
- Multinomial Coefficients
- Inclusion-Exclusion
- **Combinatorial Proofs** ◀

## Combinatorial proof: Show that $M = N$

- Let  $S$  be a set of objects
- Show how to count  $|S|$  one way  $\Rightarrow |S| = M$
- Show how to count  $|S|$  another way  $\Rightarrow |S| = N$
- Conclude that  $M = N$

## Binomial Coefficient – Many interesting and useful properties

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$$\binom{n}{n} = 1$$

$$\binom{n}{1} = n$$

$$\binom{n}{0} = 1$$

**Fact.**  $\binom{n}{k} = \binom{n}{n-k}$

Symmetry in Binomial Coefficients

**Fact.**  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

Pascal's Identity

**Fact.**  $\sum_{k=0}^n \binom{n}{k} = 2^n$

Follows from Binomial theorem

## Pascal's Identities

**Fact.**  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

How to prove Pascal's identity?

Algebraic argument:

$$\begin{aligned}\binom{n-1}{k-1} + \binom{n-1}{k} &= \frac{(n-1)!}{(k-1)!(n-k)!} + \frac{(n-1)!}{k!(n-1-k)!} \\ &= 20 \text{ years later ...} \\ &= \frac{n!}{k!(n-k)!} \\ &= \binom{n}{k}\end{aligned}$$


Hard work and not intuitive

Let's see a combinatorial argument

## Example – Binomial Identity

**Fact.**  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S| = |A| + |B|$



$S = A \cup B$ , disjoint

$S$ : the set of size  $k$  subsets of  $[n] = \{1, 2, \dots, n\} \rightarrow |S| = \binom{n}{k}$

$A$ : the set of size  $k$  subsets of  $[n]$  including  $n$


$B$ : the set of size  $k$  subsets of  $[n]$  NOT including  $n$

**Sum rule:**  
 $|A \cup B| = |A| + |B|$

## Example – Binomial Identity

**Fact.**  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S| = |A| + |B|$



**S**: the set of size  $k$  subsets of  $[n] = \{1, 2, \dots, n\}$   $\rightarrow |S| = \binom{n}{k}$

e.g.:  $n = 4$ ,  $S = \{\{1,2\}, \{1,3\}, \{1,4\}, \{2,3\}, \{2,4\}, \{3,4\}\}$

**A**: the set of size  $k$  subsets of  $[n]$  including  $n$

$$A = \{\{1,4\}, \{2,4\}, \{3,4\}\}. \quad n = 4$$

**B**: the set of size  $k$  subsets of  $[n]$  NOT including  $n$

$$B = \{\{1,2\}, \{1,3\}, \{2,3\}\}$$




## Example – Binomial Identity

**Fact.**  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

$|S|$                        $|A|$                        $|B|$

$S = A \cup B$



$S$ : the set of size  $k$  subsets of  $[n] = \{1, 2, \dots, n\}$

$A$ : the set of size  $k$  subsets of  $[n]$  including  $n$

$B$ : the set of size  $k$  subsets of  $[n]$  NOT including  $n$

$n$  is in set, need to choose  $k - 1$  elements from  $[n - 1]$

$$|A| = \binom{n-1}{k-1}$$

$n$  not in set, need to choose  $k$  elements from  $[n - 1]$

$$|B| = \binom{n-1}{k}$$

## Pascal's triangle

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$$

			1					
		1	1					
	1	2	1					
	1	3	3	1				
	1	4	6	4	1			
	1	5	10	10	5	1		
	1	6	15	20	15	6	1	
1	7	21	35	35	21	7	1	

						$\binom{0}{0}$			
					$\binom{1}{0}$	$\binom{1}{1}$			
				$\binom{2}{0}$	$\binom{2}{1}$	$\binom{2}{2}$			
			$\binom{3}{0}$	$\binom{3}{1}$	$\binom{3}{2}$	$\binom{3}{3}$			
		$\binom{4}{0}$	$\binom{4}{1}$	$\binom{4}{2}$	$\binom{4}{3}$	$\binom{4}{4}$			
$\binom{5}{0}$	$\binom{5}{1}$	$\binom{5}{2}$	$\binom{5}{3}$	$\binom{5}{4}$	$\binom{5}{5}$				

## combinatorial argument/proof

- Elegant
- Simple
- Intuitive



[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)

## Algebraic argument

- Brute force
- Less Intuitive



[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)

## Combinatorial Proof?

**Fact.**  $\sum_{k=0}^n \binom{n}{k} = 2^n$

