


CSE 421 Introduction to Algorithms Winter 2000

The Network Flow Problem

CSE 421,W '00, Ruzzo

1

The Network Flow Problem



- How much stuff can flow from s to t ?

CSE 421,W '00, Ruzzo

2

Net Flow: Formal Definition

Given:

A digraph $G = (V, E)$

Two vertices s, t in V
(source & sink)

A capacity $c(u, v) \geq 0$
for each $(u, v) \in E$
(and $c(u, v) = 0$ for all non-edges (u, v))

Find:

A flow function $f: V \times V \rightarrow \mathbb{R}$ s.t.,
for all u, v :

- $f(u, v) \leq c(u, v)$ [Capacity Constraint]
- $f(u, v) = -f(v, u)$ [Skew Symmetry]
- if $u \neq s, t$, $f(u, V) = 0$ [Flow Conservation]

Maximizing total flow $|f| = f(s, V)$


Notation:

$$f(X, Y) = \sum_{x \in X} \sum_{y \in Y} f(x, y)$$

CSE 421,W '00, Ruzzo

3

Example: A Flow Function



$$f(s, u) = f(u, t) = 2$$


$$f(u, s) = f(t, u) = -2$$

$$f(u, V) = \sum_{v \in V} f(u, v) = f(u, s) + f(u, t) = -2 + 2 = 0$$

CSE 421,W '00, Ruzzo

4

Example: A Flow Function



- Not shown: $f(u, v)$ if ≤ 0
- Note: max flow ≥ 4 since f is a flow function, with $|f| = 4$

CSE 421,W '00, Ruzzo

5

Max Flow via a Greedy Alg?

While there is an $s \rightarrow t$ path in G


Pick such a path, p

Find c , the min capacity of any edge in p

Subtract c from all capacities on p

Delete edges of capacity 0

- This does NOT always find a max flow:



If pick $s \rightarrow b \rightarrow a \rightarrow t$ first, flow stuck at 2.
But flow 3 possible.

CSE 421,W '00, Ruzzo

6

A Brief History of Flow


#	year	discoverer(s)	bound
1	1951	Dinic	$O(n^m U)$
2	1955	Ford & Fulkerson	$O(n m U)$
3	1970	Dinits	$O(n m^2)$
4	1971		$O(n^2 m)$
5	1972	Edmonds & Karp Dinits	$O(m n \log U)$
6	1973	Dinits	$O(n m \log U)$
7	1974	Gabow	$O(n^2)$
8	1977	Cherkassky	$O(n^2 \sqrt{m})$
9	1980	Gull & Naamad	$O(n m \log^2 n)$
10	1983	Borod & Tarjan	$O(n m \log n)$
11	1984	Goldberg & Tarjan	$O(n m \log(n/m))$
12	1987	Aluja & Orlin	$O(n m + n^2 \log U)$
13	1987	Ahuja et al.	$O(n m \log(n \log U)/(m+2))$
14	1989	Cheiryan & Hagerup	$O(n^2) \log n$
15	1990	Alon & Cheiryan et al.	$O(n^2) \log^2 n$
16	1990	Alon	$O(n m + n^{2/3} \log n)$
17	1992	King et al.	$O(n m + n^{2/3})$
18	1993	Phillips & Westbrook	$O(n m (\log_{n/(m+n)} n + \log^{2/3} n))$
19	1994	King et al.	$O(n m \log_{n/(m+n)} n)$
20	1997	Goldberg & Rao	$O(n^{2/3} m \log(n/m) \log U)$
			$O(n^{2/3} m \log(n/m) \log U)$

n = # of vertices
m = # of edges
U = Max capacity

CSE 421, W '00, Ruzzo

7

Greed Revisited




CSE 421, W '00, Ruzzo

8

Residual Capacity

- The *residual capacity* (w.r.t. f) of (u,v) is $c_f(u,v) = c(u,v) - f(u,v)$



- e.g. $c_f(s,b)=7$; $c_f(a,x) = 1$; $c_f(x,a) = 3$

CSE 421, W '00, Ruzzo

9

Residual Networks & Augmenting Paths

- The *residual network* (w.r.t. f) is the graph $G_f = (V, E_f)$, where


$$E_f = \{ (u,v) \mid c_f(u,v) > 0 \}$$

- An *augmenting path* (w.r.t. f) is a simple $s \rightarrow t$ path in G_f .

CSE 421, W '00, Ruzzo


10

A Residual Network



11

An Augmenting Path



12

Lemma 1


If f admits an augmenting path p , then f is not maximal.

Proof: "obvious" -- augment along p by c_p , the min residual capacity of p 's edges.

CSE 421, W '00, Ruzzo

13


Augmenting A Flow



CSE 421, W '00, Ruzzo

14


Proof of Lemma 1



CSE 421, W '00, Ruzzo

15

Proof of Lemma 1



16

Ford-Fulkerson Method

While G_f has an augmenting path, augment

- Questions:


- » Does it halt?
- » Does it find a maximum flow?
- » How fast?

CSE 421, W '00, Ruzzo

17

Cuts

- A partition S, T of V is a **cut** if $s \in S, t \in T$
- Capacity** of cut S, T is $c(S, T) = \sum_{u \in S} \sum_{v \in T} c(u, v)$




CSE 421, W '00, Ruzzo

18

Lemma 2

- For any flow f and any cut S, T ,
 - the net flow across the cut equals the total flow, i.e., $|f| = f(S, T)$, and
 - the net flow across the cut cannot exceed the capacity of the cut, i.e. $f(S, T) \leq c(S, T)$
- Corollary:**
Max flow \leq Min cut



CSE 421, W '00, Ruzzo

19

Max Flow / Min Cut Theorem

- For any flow f , the following are equivalent
- $|f| = c(S, T)$ for some cut S, T (a min cut)
 - f is a maximum flow
 - f admits no augmenting path


Proof:

- $(1) \Rightarrow (2)$: corollary to lemma 2
- $(2) \Rightarrow (3)$: lemma 1

CSE 421, W '00, Ruzzo

20

(3) \Rightarrow (1)



$S = \{ u \mid \exists \text{ an augmenting path from } s \text{ to } u \}$
 $T = V - S; s \in S, t \in T$

For any (u, v) in $S \times T$, \exists an augmenting path from s to u , but **not** to v .

$\therefore (u, v)$ has 0 residual capacity:

$$(u, v) \in E \Rightarrow \text{saturated} \quad f(u, v) = c(u, v)$$

$$(v, u) \in E \Rightarrow \text{no flow} \quad f(u, v) = f(v, u) = 0$$

This is true for every edge crossing the cut, i.e.


$$|f| = f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) = \sum_{u \in S, v \in T, (u, v) \in E} f(u, v) = \sum_{u \in S, v \in T, (u, v) \in E} c(u, v) = c(S, T)$$

CSE 421, W '00, Ruzzo

21

Corollaries & Facts

- If Ford-Fulkerson terminates, then it's found a max flow.
- It will terminate if $c(e)$ integer or rational (but may not if they're irrational).
- However, may take exponential time, even with integer capacities:



$c = 10^9$, say

CSE 421, W '00, Ruzzo

22

Edmonds-Karp Algorithm

- Use a **shortest** augmenting path (via Breadth First Search in residual graph)
- Time: $O(n m^2)$


CSE 421, W '00, Ruzzo

23

BFS/Shortest Path Lemmas

Distance from s is never reduced by:

- Deleting** an edge
proof: no new (hence no shorter) path created
- Adding** an edge (u, v) , **provided** v is nearer than u
proof: BFS is unchanged, since v visited before (u, v) examined



CSE 421, W '00, Ruzzo

24

Lemma 27.8 (Alternate Proof)


Let f be a flow, G_f the residual graph, and p a shortest augmenting path. Then no vertex is closer to s after augmentation along p .

Proof: Augmentation only deletes edges, adds back edges

CSE 421, W '00, Ruzzo

25

Augmentation vs BFS



CSE 421, W '00, Ruzzo

26

Theorem 27.9

The Edmonds-Karp Algorithm performs $O(mn)$ flow augmentations

Proof:

$\{u,v\}$ is **critical** on augmenting path p if it's closest to s having min residual capacity won't be critical again until farther from s so each edge critical at most n times

CSE 421, W '00, Ruzzo

27

Corollary

- Edmonds-Karp runs in $O(nm^2)$


CSE 421, W '00, Ruzzo

28

Flow Integrality Theorem

If all capacities are integers

- The max flow has an integer value
- Ford-Fulkerson method finds a max flow in which $f(u,v)$ is an integer for all edges (u,v)



CSE 421, W '00, Ruzzo

29