

## 1 Triangles in Graphs (Optional)

**Theorem 1.** *If a graph on  $2n$  vertices has  $n^2 + 1$  edges, then it has a triangle.*

**Proof** We prove it by induction on  $n$ .  $P(n)$  = “Any graph  $G = (V, E)$  with  $2n$  vertices and  $m \geq n^2 + 1$  edges has a triangle.

Base Case:  $P(1)$ : When  $n = 1$ ,  $P(1)$  holds since the number of edges is at most  $1 < n^2 + 1$ .

IH:  $P(n)$  holds for some  $n \geq 1$ .

IS: We prove  $P(n + 1)$ . Let  $G$  be an arbitrary graph with  $2(n + 1)$  vertices and at least  $m \geq (n + 1)^2 + 1$  edges. Let  $\{x, y\}$  be an arbitrary edge in the graph. Consider the graph  $G' = G - x - y$  on  $2n$  vertices obtained by **deleting**  $x, y$  (and all of their incident edges) from the original graph. If  $G'$  has at least  $n^2 + 1$  edges, then by IH it has a triangle, and we are done.

Otherwise,  $G'$  has at most  $n^2$  edges. Since  $G$  has at least  $(n + 1)^2 + 1$  edges, by removing  $x, y$  we have deleted  $(n + 1)^2 + 1 - n^2 = 2n + 2$  edges from  $G$ . Since  $\{x, y\}$  is also an edge, there are at least  $2n + 1$  edges that connect  $x, y$  to the vertices of  $G'$ . Thus by the pigeonhole principle, there is some vertex  $z$  so that  $\{x, z\}, \{y, z\}$  are both edges of  $G$ . But, then  $x, y, z$  form a triangle in  $G$ . ■

The above theorem is tight. Consider the graph with  $n$  vertices on the left and  $n$  vertices on the right and every vertex on the left is connected to every vertex on the right. This graph has no triangles but  $n^2$  edges.

Also, note the importance of deletion in the induction. Here, we crucially used that the  $x, y$  pair deleted from  $G$  were **neighbors** in  $G$ .