
CSE 331

Software Design & Implementation

Section 3 – HW4, Abstract Data Types, and JUnit

Agenda

- Overview of HW4
- Quick review of polynomial arithmetic
- Review abstract data types (ADTs) by example
- Unit testing with Junit – an initial tour for HW4

HW4 – Polynomial calculator

A homework in 6 parts:

0. Pseudocode algorithms for polynomial arithmetic
1. Conceptual questions about **RatNum**
2. Implement **RatTerm**
3. Implement **RatPoly**
4. Implement **RatPolyStack**
5. Try out your finished calculator!
6. Run your code against our tests to make sure it works!

Start early, and use your knowledge of invariants to unblock yourself.




The RatThings


- **RatNum** ADT
 - A rational number
 - Also includes a NaN (“not a number”) value
- **RatTerm** ADT
 - A polynomial term (rational coefficient w/ integer degree)
- **RatPoly** ADT
 - A polynomial expression (sum of polynomial terms)
- **RatPolyStack** ADT
 - An ordered collection of polynomial expressions



The RatThings



The RatThings



Polynomial arithmetic

Review arithmetic operations over polynomial expressions:

1. Addition
2. Subtraction
3. Multiplication
4. Division

Defining and following invariants is critical to making sure that these operations are implemented correctly.

Polynomial addition

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$

Polynomial addition

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 5x^4 + 4x^3 - 1x^2 + 5 \\ + 3x^5 - 2x^3 + 1x - 5 \\ \hline \end{array}$$

Polynomial addition

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 \ 5 \ +x^4 \ 4 \ +x^3 \ - \ 1x^2 \ + \ 0x \ + \ 5 \\ + \quad 3x^5 + \ 0x^4 \ - \ 2x^3 \ + \ 0x^2 \ + \ 1x \ - \ 5 \\ \hline \end{array}$$

Polynomial addition

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 + 5x^4 + 4x^3 - 1x^2 + 0x + 5 \\ + \quad 3x^5 + 0x^4 - 2x^3 + 0x^2 + 1x - 5 \\ \hline 3x^5 + 5x^4 + 2x^3 - 1x^2 + 1x + 0 \end{array}$$

Polynomial subtraction

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

Polynomial subtraction

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 5x^4 + 4x^3 - 1x^2 + 5 \\ - 3x^5 - 2x^3 + 1x - 5 \\ \hline \end{array}$$

Polynomial subtraction

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 + 5x^4 + 4x^3 - 1x^2 + 0x + 5 \\ - 3x^5 + 0x^4 - 2x^3 + 0x^2 + 1x - 5 \\ \hline \end{array}$$

Polynomial subtraction

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 + 5x^4 + 4x^3 - 1x^2 + 0x + 5 \\ - 3x^5 + 0x^4 - 2x^3 + 0x^2 + 1x - 5 \\ \hline -3x^5 + 5x^4 + 6x^3 - 1x^2 - 1x + 10 \end{array}$$

Polynomial multiplication

$$(4x^3 - x^2 + 5) \times (x - 5)$$

Polynomial multiplication

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} 4x^3 - 1x^2 + 5 \\ \times \qquad \qquad \qquad 1x - 5 \\ \hline \end{array}$$

Polynomial multiplication

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} 4x^3 - 1x^2 \\ \times \qquad \qquad \qquad 1x \\ \hline -20x^3 + 5x^2 \qquad \qquad \qquad + 5 \\ \qquad \qquad \qquad - 5 \end{array}$$

The diagram shows the multiplication of two polynomials using the vertical method. The first polynomial, $4x^3 - x^2 + 5$, is written above the multiplication sign. The second polynomial, $x - 5$, is written to the right. A horizontal line separates the terms of the first polynomial from the product. The term -5 in the second polynomial is circled in green.

Polynomial multiplication

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} 4x^3 - 1x^2 + 5 \\ \times \quad \quad \quad 1x - 5 \\ \hline -20x^3 + 5x^2 \qquad \qquad \qquad - 25 \\ 4x^4 - 1x^3 \qquad \qquad \qquad + 5x \end{array}$$

Polynomial multiplication

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} 4x^3 - 1x^2 + 5 \\ \times \quad \quad \quad 1x - 5 \\ \hline -20x^3 + 5x^2 \quad \quad \quad - 25 \\ + \quad 4x^4 - 1x^3 \quad \quad \quad + 5x \\ \hline 4x^4 - 21x^3 + 5x^2 + 5x - 25 \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 1x^3 \\ -2x \quad -5 \end{array} \overline{\left[\begin{array}{r} 5x^6 \\ +4x^4 \\ -1x^3 \\ +5 \end{array} \right]}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 1x^3 + 0x^2 - 2x - 5 \quad \hline 5x^6 \quad +0x^5 \quad +4x^4 \quad -1x^3 \quad +0x^2 \quad +0x \quad +5 \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 \\ \hline 1x^3 + 0x^2 - 2x - 5 & \overline{)5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 \\ \hline 1x^3 + 0x^2 - 2x - 5 & \overline{)5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ & 5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5 \\ & -5x^6 & -0x^5 & -10x^4 & -25x^3 & & & \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 \\ \hline 1x^3 + 0x^2 - 2x - 5 & \overline{)5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ & - 5x^6 + 0x^5 - 10x^4 - 25x^3 \\ \hline & 0x^6 + 0x^5 + 14x^4 + 24x^3 \end{array}$$

Notice (quotient * divisor) + remainder is always equal to $(5x^6 + 4x^4 - x^3 + 5)$.

We can use this fact to produce an invariant.

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 \\ \hline 1x^3 + 0x^2 - 2x - 5 & \overline{)5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ & - 5x^6 + 0x^5 - 10x^4 - 25x^3 \\ \hline & 0x^6 + 0x^5 + 14x^4 + 24x^3 \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 \\ \hline 1x^3 + 0x^2 - 2x - 5 & \overline{)5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ & - 5x^6 + 0x^5 - 10x^4 - 25x^3 \\ \hline & 0x^6 + 0x^5 + 14x^4 + 24x^3 + 0x^2 \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 \\ \hline 1x^3 +0x^2 & -2x & -5 & \left[\begin{array}{rrrrrr} 5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5 \\ - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & & \end{array} \right] \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 \\ \hline 1x^3 +0x^2 & -2x & -5 & | & 5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5 \\ & - & 5x^6 & +0x^5 & -10x^4 & -25x^3 \\ \hline & & 0x^6 & +0x^5 & +14x^4 & +24x^3 & & +0x^2 & +0x \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x \\ \hline 1x^3 +0x^2 -2x -5 & \overline{)5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5} \\ & - & 5x^6 & +0x^5 & -10x^4 & -25x^3 \\ \hline & & 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x \\ \hline 1x^3 +0x^2 & -2x & -5 & \left[\begin{array}{rrrrrr} 5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5 \\ - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x & \\ & & 14x^4 & +0x^3 & -28x^2 & -70x & \end{array} \right] \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x \\ \hline 1x^3 +0x^2 & -2x & -5 & \left[\begin{array}{rrrrrr} 5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5 \\ - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x & \\ - & & 14x^4 & +0x^3 & -28x^2 & -70x & \\ \hline 0x^4 & +24x^3 & +28x^2 & +70x & & & \end{array} \right] \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x \\ \hline 1x^3 +0x^2 -2x -5 & \overline{)5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5} \\ & - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline & & 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x \\ & & - & 14x^4 & +0x^3 & -28x^2 & -70x & \\ \hline & & 0x^4 & +24x^3 & +28x^2 & +70x & & \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x \\ \hline 1x^3 +0x^2 -2x -5 & \overline{)5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5} \\ & - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline & & 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x \\ & & - & & 14x^4 & +0x^3 & -28x^2 & -70x \\ \hline & & & & 0x^4 & +24x^3 & +28x^2 & +70x & +5 \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x & +24 \\ \hline 1x^3 +0x^2 & -2x & -5 & \left[\begin{array}{rrrrrr} 5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5 \\ - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x & \\ - & & 14x^4 & +0x^3 & -28x^2 & -70x & \\ \hline 0x^4 & +24x^3 & +28x^2 & +70x & & & +5 \end{array} \right] \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x & +24 \\ \hline 1x^3 +0x^2 -2x -5 & \overline{)5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5} \\ & - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline & & 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x & \\ & & - & 14x^4 & +0x^3 & -28x^2 & -70x & \\ \hline & & 0x^4 & +24x^3 & +28x^2 & +70x & +5 \\ & & & 24x^3 & +0x^2 & -48x & -120 \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} & & 5x^3 & +0x^2 & +14x & +24 \\ \hline 1x^3 +0x^2 -2x -5 & \overline{)5x^6 & +0x^5 & +4x^4 & -1x^3 & +0x^2 & +0x & +5} \\ & - & 5x^6 & +0x^5 & -10x^4 & -25x^3 & & \\ \hline & & 0x^6 & +0x^5 & +14x^4 & +24x^3 & +0x^2 & +0x & \\ & & - & 14x^4 & +0x^3 & -28x^2 & -70x & & \\ \hline & & 0x^4 & +24x^3 & +28x^2 & +70x & +5 & & \\ & & - & 24x^3 & +0x^2 & -48x & -120 & & \\ \hline & & 0x^3 & +28x^2 & +118x & +125 & & & \end{array}$$

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

quotient

$$\begin{array}{r} 5x^3 \quad +0x^2 \quad +14x \quad +24 \\ \hline 0x^6 \quad +0x^5 \quad +14x^4 \quad +24x^3 \quad +0x^2 \quad +0x \quad +5 \\ - 5x^6 \quad +0x^5 \quad -10x^4 \quad -25x^3 \\ \hline 0x^6 \quad +0x^5 \quad +14x^4 \quad +24x^3 \quad +0x^2 \quad +0x \\ - 14x^4 \quad +0x^3 \quad -28x^2 \quad -70x \\ \hline 0x^4 \quad +24x^3 \quad +28x^2 \quad +70x \quad +5 \\ - 24x^3 \quad +0x^2 \quad -48x \quad -120 \\ \hline 0x^3 \quad +28x^2 \quad +118x \quad +125 \end{array}$$

remainder

Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$5x^3 + 14x + 24 + \frac{28x^2 + 118x + 125}{x^3 - 2x - 5}$$

Notice that the loop invariant, $q*y + r = x$ and $0 \leq r$ where q is the quotient, y is the divisor, r is the remainder and x is the polynomial that is being divided is always correct after each subtraction step.

Abstract data types by example

Review ADT concepts through two examples:


- A **Line** ADT
- A **Rectangle** ADT

On the course website, see “Resources” → “Class and Method Specifications” for a handy guide with full details.

We won’t cover representation invariants today (see Friday’s lecture).


Line ADT

Concept: A line segment in the Cartesian co-ordinate plane



Line ADT: Representation #1


```
/**  
 * A Line is a mutable 2D line segment with endpoints  
 * p1 and p2.  
 */  
  
public class Line {  
    // Abstract state is line with endpoints p1 and p2  
    private Point p1, p2;  
}
```



Line ADT: Representation #2

```
/**  
 * A Line is a mutable 2D line segment with endpoints  
 * p1 and p2.  
 */  
  
public class Line {  
    // Abstract state is line with endpoints (x1, y1) and  
    //                                         (x2, y2)  
    private int x1, x2;  
    private int y1, y2;  
}
```


Does this representation have any advantages over #1?



Line ADT: Representation #3

```
/**  
 * A Line is a mutable 2D line segment with endpoints  
 * p1 and p2.  
 */  
  
public class Line {  
    // Abstract state is line with endpoints (x1, y1) and  
    // (x1 + len * cos(angle), y1 + len * sin(angle))  
    private int x1, y1;  
    private double angle;  
    private double len;  
}
```

Does this representation have any advantages over #1?



Try it yourself!

Write your own specification of a Rectangle ADT on the handout.

Then give two different possible representations for your Rectangle ADT and write abstraction functions for them

Testing: A quick introduction

- For HW 4, you'll be running our test suite to verify your RatThings work.
- Let's do a quick walkthrough of our test suite
 - Just know how it works; don't need to know how to write tests!

JUnit

- Industry-standard Java toolkit for unit testing
 - We're using JUnit 4
- A unit test is a test for one “component” by itself
 - “Component” typically a class or a method
- Each unit test written as a method
 - We'll see the particulars in a moment...
- Closely related unit tests should be grouped into a class
 - For example, all unit tests for the same ADT implementation

Writing tests with JUnit

Annotate a method with `@Test` to flag it as a JUnit test

```
import org.junit.*;
import static org.junit.Assert.*;

/** Unit tests for my Foo ADT implementation */
public class FooTests {
    @Test
    public void testBar() {
        ... /* use JUnit assertions in here */
    }
}
```

Using JUnit assertions

- JUnit assertions establish success or failure of the test method
 - *Note: JUnit assertions are *different* from Java's `assert` statement*
- Use to check that an actual result matches the expected value
 - Example: `assertEquals(42, meaningOfLife());`
 - Example: `assertTrue(list.isEmpty());`
- A test method stops immediately after the first assertion failure
 - If no assertion fails, then the test method passes
 - Other test methods still run either way
- JUnit results show details of any test failures

Common JUnit assertions

JUnit's documentation has a full list, but these are the most common assertions.

Assertion	Failure condition
<code>assertTrue(test)</code>	<code>test == false</code>
<code>assertFalse(test)</code>	<code>test == true</code>
<code>assertEquals(expected, actual)</code>	<code>expected</code> and <code>actual</code> are not equal
<code>assertSame(expected, actual)</code>	<code>expected != actual</code>
<code>assertNotSame(expected, actual)</code>	<code>expected == actual</code>
<code>assertNull(value)</code>	<code>value != null</code>
<code>assertNotNull(value)</code>	<code>value == null</code>

Any JUnit assertion can also take a string to show in case of failure, e.g.,
`assertEquals("helpful message", expected, actual)`.

Checking for a thrown exception

- Should test that your code throws exceptions as specified
- This kind of test method fails if its body does *not* throw an exception of the named class
 - May not need any JUnit assertions inside the test method unlike our previous guideline

```
@Test(expected=IndexOutOfBoundsException.class)
public void testGetEmptyList() {
    List<String> list = new ArrayList<String>();
    list.get(0);
}
```

Test ordering, setup, clean-up

JUnit does not promise to run tests in any particular order.

However, JUnit can run helper methods for common setup/cleanup

- Run before/after each test method in the class:

```
@Before  
public void m() { ... }  
  
@After  
public void m() { ... }
```

- Run once before/after running *all* test methods in the class:

```
@BeforeClass  
public static void m() { ... }  
  
@AfterClass  
public static void m() { ... }
```