Particle Dynamics

Reading

Particle Systems Dynamics handout

Optional:

Hocknew and Eastwood. *Computer simulation using particles*. Adam Hilger, New York, 1988.

Gavin Miller. "The motion dynamics of snakes and worms." *Computer Graphics* 22:169-178, 1988.

2

Overview

- One lousy particle
- Particle systems
- Forces: gravity, springs
- Implementation

3

Newtonian particle

- Differential equations: f=ma
- Forces depend on:
- Position, velocity, time

$$\ddot{x} = \frac{f(x, \dot{x}, t)}{m}$$

4

Second order equations

$$\ddot{x} = \frac{f(x, \dot{x}, t)}{m}$$

Has 2nd derivatives

$$\begin{bmatrix} \dot{x} = v \\ \dot{v} = \frac{f(x, \dot{x}, t)}{m} \end{bmatrix}$$

Add a new variable v to get
a pair of coupled 1st order equations

5

Phase space

$$\begin{bmatrix} x \\ v \end{bmatrix}$$

Concatenate x and v to make a 6-vector:
position in phase space

$$\begin{bmatrix} \dot{x} \\ \dot{v} \end{bmatrix}$$

Velocity on Phase space:
Another 6-vector

$$\begin{bmatrix} \dot{x} \\ \dot{v} \end{bmatrix} = \begin{bmatrix} v \\ f / m \end{bmatrix}$$

A vanilla 1st-order differential equation

6

Particle structure

Position in phase space

$$\begin{bmatrix} \mathbf{x} \\ \mathbf{v} \\ \mathbf{f} \\ m \end{bmatrix}$$

- $\mathbf{x}$ ← position
- $\mathbf{v}$ ← velocity
- $\mathbf{f}$ ← force accumulator
- m ← mass

7

Solver interface



8

Particle systems



9

Solver interface



10

Differential equation solver

$$\begin{bmatrix} \dot{x} \\ \dot{v} \end{bmatrix} = \begin{bmatrix} v \\ f/m \end{bmatrix}$$

Euler method: $x(t+h) = x(t) + h \cdot \dot{x}(t)$

$$\mathbf{x}_{i+1} = \mathbf{x}_i + \Delta t \cdot \dot{\mathbf{x}}_i$$

$$\mathbf{v}_{i+1} = \mathbf{v}_i + \Delta t \cdot \dot{\mathbf{v}}_i$$

Gets very unstable for large Δt

Higher order solvers perform better: (e.g. Runge-Kutta)

11

derivEval loop

1. Clear forces
   - Loop over particles, zero force accumulators
2. Calculate forces
   - Sum all forces into accumulators
3. Gather
   - Loop over particles, copying v and f/m into destination array

12

Forces

- Constant (gravity)
- Position/time dependent (force fields)
- Velocity-dependent (drag)
- N-ary (springs)

13

Force structures

Force objects are black boxes that point to the particles they influence, and add in their contribution into the force accumulator.

Global force calculation:

- Loop, invoking force objects

14

Particle systems with forces



15

Gravity

Force law:

$\mathbf{f}_{grav} = m\mathbf{G}$

```
p->f += p->m * F->G
```

16

Viscous drag

Force law:

$\mathbf{f}_{drag} = -k_{drag}\mathbf{v}$

```
p->f -= F->k * p->v
```

17

Damped spring

Force law:

$$\mathbf{f}_1 = -\left[k_s(|\mathrm{V}\mathbf{x}| - \mathbf{r}) + k_d\left(\frac{\mathrm{V}\mathbf{v}\mathrm{V}\mathbf{x}}{|\mathrm{V}\mathbf{x}|}\right)\right]\frac{\mathrm{V}\mathbf{x}}{|\mathrm{V}\mathbf{x}|}$$

$\mathbf{f}_2 = -\mathbf{f}_1$



18

derivEval Loop



19

Solver interface



20

Differential equation solver

$$\begin{bmatrix} \dot{x} \\ \dot{v} \end{bmatrix} = \begin{bmatrix} v \\ f/m \end{bmatrix}$$

Euler method:

$$\begin{bmatrix} x_1^{i+1} \\ v_1^{i+1} \\ \mathrm{M} \\ x_n^{i+1} \\ v_n^{i+1} \end{bmatrix} = \begin{bmatrix} x_1^{i} \\ v_1^{i} \\ \mathrm{M} \\ x_n^{i} \\ v_n^{i} \end{bmatrix} + \Delta t \begin{bmatrix} v_1^{i} \\ f_1^{i}/m_1 \\ \mathrm{M} \\ v_n^{i} \\ f_n^{i}/m_n \end{bmatrix}$$

21

Bouncing off the walls



- Add-on for a particle simulator
- For now, just simple point-plane collisions

22

Normal and tangential components





$$V_N = (N \cdot V)N$$
$$V_T = V - V_N$$

23

Collision Detection



$(\mathbf{X}-\mathbf{P})\cdot\mathbf{N} < \varepsilon$ Within e of the wall

$\mathbf{N}\cdot\mathbf{V} < 0$ Heading in

24

Collision Response



before after

$$\mathbf{V}' = \mathbf{V}_T - k_r\mathbf{V}_N$$

25

Summary

- Physics of a particle system
- Various forces acting on a particle
- Combining particles into a particle system
- Euler method for solving differential equations

26