

CSE 312

Foundations of Computing II

Lecture 2: Combinations and Binomial Coefficients

Announcements

Office hours start today:

- In particular, I will be available for office hours starting right after class today (CSE 668)

Homework:

- Pset1 was posted on Wednesday and is due 11:59pm next Wednesday.
- We will have the same pattern for the other assignments, excluding the week of the midterm and the last two assignments after Thanksgiving, which will be due on Fridays.

Python programming on homework:

- Some problem sets will include coding problems
 - in Python (*no prior knowledge or experience required*)
 - provide a deeper understanding of how theory we discuss is used in practice
 - should be fun

Announcements

- **Problem Set 1**

- Read the first page for how to write up your homework solutions. Don't wait until you are working on the questions to figure it out!
- Section solutions are a good place to look at for examples.

- **Resources**

- Textbook readings can provide another perspective
- **Theorems & Definitions sheet** – https://www.alextsun.com/files/defs_thms.pdf
- Office Hours
- EdStem discussion

- **EdStem discussion etiquette**

- OK to publicly discuss content of the course and any confusion over topics discussed in class, but **not solutions** for current homework problems, or anything about current exams that have not yet been graded.
- It is also acceptable to ask for clarifications about what current homework problems are asking and concepts behind them, just not about their solutions.

Quick counting summary from last class

- **Sum rule:**

If you can choose from

- EITHER one of n options,
- OR one of m options with NO overlap with the previous n ,

then the number of possible outcomes of the experiment is $n + m$

- **Product rule:**

In a sequential process, if there are

- n_1 choices for the 1st step,
- n_2 choices for the 2nd step (given the first choice), ..., and
- n_k choices for the k^{th} step (given the previous choices),

then the total number of outcomes is $n_1 \times n_2 \times n_3 \times \cdots \times n_k$

- Representation of the problem is important (creative part)

Factorial

“How many ways to order elements in S , where $|S| = n$?”

Permutations

$$\text{Answer} = n \times (n - 1) \times (n - 2) \times \cdots \times 2 \times 1$$

Definition. The factorial function is

$$n! = n \times (n - 1) \times \cdots \times 2 \times 1$$

Note: $0! = 1$

Theorem. (Stirling's approximation)

$$\underbrace{\sqrt{2\pi}}_{= 2.5066} \cdot n^{n+\frac{1}{2}} \cdot e^{-n} \leq n! \leq \underbrace{e}_{= 2.7183} \cdot n^{n+\frac{1}{2}} \cdot e^{-n}$$

Huge: Grows exponentially in n

Distinct Letters

*“How many sequences of 5 **distinct** alphabet letters from $\{A, B, \dots, Z\}$?”*

E.g., AZURE, BINGO, TANGO. But not: STEVE, SARAH

Answer: $26 \times 25 \times 24 \times 23 \times 22$
 $= 7893600$

In general



Aka: k -permutations

Fact. # of k -element sequences of distinct symbols from an n -element set is

$$P(n, k) = n \times (n - 1) \times \cdots \times (n - k + 1) = \frac{n!}{(n - k)!}$$

Today: More Counting

- Permutations and Combinations



Number of Subsets

*“How many size-5 **subsets** of $\{A, B, \dots, Z\}$?”*

E.g., $\{A, Z, U, R, E\}$, $\{B, I, N, G, O\}$, $\{T, A, N, G, O\}$. But not:
 $\{S, T, E, V\}$, $\{S, A, R, H\}$, ...

Difference from k -permutations: NO ORDER

Different sequences: TANGO, OGNAT, ATNGO, NATGO, ONATG ...

Same set: $\{T, A, N, G, O\}$, $\{O, G, N, A, T\}$, $\{A, T, N, G, O\}$, $\{N, A, T, G, O\}$, $\{O, N, A, T, G\}$

Number of Subsets – Idea

Consider a sequential process:

1. Choose a subset $S \subseteq \{A, B, \dots, Z\}$ of size $|S| = 5$
e.g. $S = \{A, G, N, O, T\}$
2. Choose a permutation of letters in S
e.g., *TANGO, AGNOT, NAGOT, GOTAN, GOATN, NGOAT, ...*

Outcome: A sequence of 5 distinct letters from $\{A, B, \dots, Z\}$

$$??? = \frac{26!}{21! 5!} = 65780$$

$$\begin{array}{c} \boxed{???} \\ \times \\ \boxed{5!} \\ = \\ \boxed{\frac{26!}{21!}} \end{array}$$

Number of Subsets – Binomial Coefficient

Fact. The number of subsets of size k of a set of size n is

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

Binomial coefficient (verbalized as “ n choose k ”)

Notation: $\binom{S}{k}$ = set of all k -element subsets of S . $\left| \binom{S}{k} \right| = \binom{|S|}{k}$
[also called **combinations**]

Symmetry in Binomial Coefficients

Fact. $\binom{n}{k} = \binom{n}{n-k}$

This is called an Algebraic proof,
i.e., Prove by checking algebra

Proof. $\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n!}{(n-k)!k!} = \binom{n}{n-k}$

Why??




Symmetry in Binomial Coefficients – A different proof


Fact. $\binom{n}{k} = \binom{n}{n-k}$

Two **equivalent** ways to choose k out of n objects (unordered)


1. Choose which k elements are **included**
2. Choose which $n - k$ elements are **excluded**



$$\binom{4}{1} = 4 = \binom{4}{3}$$




Example – Counting Paths




“How many shortest paths from Gates to Starbucks?”

Example – Counting Paths



How do we represent a shortest path?

Example – Counting Paths



Path $\in \{\uparrow, \rightarrow\}^7$

$(\uparrow, \rightarrow, \uparrow, \rightarrow, \rightarrow, \uparrow, \rightarrow)$

\uparrow 's = 3, # \rightarrow 's = 4

Poll: # of shortest paths?

A. 2^7

B. $\frac{7!}{4!}$

C. $\binom{7}{4} = \frac{7!}{4!3!}$

D. No idea

<https://pollev.com/paulbeameo28>



Example – Sum of integers

“How many solutions (x_1, \dots, x_k) such that $x_1, \dots, x_k \geq 0$ and $\sum_{i=1}^k x_i = n$?”

Example: $k = 3, n = 5$

$(0,0,5), (5,0,0), (1,0,4), (2,1,2), (3,1,1), (2,3,0), \dots$

Hint: we can represent each solution as a binary string.

Example – Sum of integers

Example: $k = 3, n = 5$

$(0,0,5), (5,0,0), (1,0,4), (2,1,2), (3,1,1), (2,3,0), \dots$

Clever representation of solutions


$(3,1,1)$	$(2,1,2)$	$(1,0,4)$
↓	↓	↓
1 1 1 0 1 0 1	1 1 0 1 0 1 1	1 0 0 1 1 1 1

Example – Sum of integers

Example: $k = 3, n = 5$

sols = # strings from $\{0,1\}^7$ w/ exactly two 0s = $\binom{7}{2} = 21$

Clever representation of solutions



Example – Sum of integers

“How many solutions (x_1, \dots, x_k) such that $x_1, \dots, x_k \geq 0$ and $\sum_{i=1}^k x_i = n$?”

$$\begin{aligned} \# \text{ sols} &= \# \text{ strings from } \{0,1\}^{n+k-1} \text{ w/ } k-1 \text{ 0s} \\ &= \binom{n+k-1}{k-1} \end{aligned}$$


After a change in representation, the problem magically reduces to counting combinations.

Example – Sum of integers

Example: $k = 3, n = 5$

sols = # strings from $\{0,1\}^7$ w/ exactly two 0s = $\binom{7}{2} = 21$

Clever representation of solutions



Also called the “stars and bars” method

*aka. “stars and bars method”

More general counting using binary encoding*

The number of ways to distribute n indistinguishable balls into k distinguishable bins is

$$\binom{n + k - 1}{k - 1} = \binom{n + k - 1}{n}$$

Example with k non-negative integers summing to n :
bins are the k integers, balls are the n 1's that add to n .

Coins

How many ways can you distribute 32 identical coins among Alex, Barbara, Charlie, Dana, and Eve?

1. Identify balls
2. Identify bins

$$\binom{32 + 5 - 1}{5 - 1}$$



A mixed example – Word Permutations (aka Anagrams)

“How many ways to re-arrange the letters in the word SEATTLE?”

STALEET, TEALEST, LASTTEE, ...

Guess: 7! Correct?!


No! e.g., swapping two T’s also leads to *SEATTLE*
swapping two E’s also leads to *SEATTLE*

Counted as separate permutations, but they lead to the same word.

A mixed example – Word Permutations (aka Anagrams)

“How many ways to re-arrange the letters in the word SEATTLE?”

STALEET, TEALEST, LASTTEE, ...



Another way to look at SEATTLE

“How many ways to re-arrange the letters in the word SEATTLE?”

STALEET, TEALEST, LASTTEE, ...

$$\binom{7}{2} \times \binom{5}{2} \times 3 \times 2 \times 1 = \frac{7!}{2! \cancel{5!}} \times \frac{\cancel{5!}}{2! \cancel{3!}} \times \cancel{3!}$$
$$= \frac{7!}{2! 2!} = 1260$$

Another interpretation:

Arrange the 7 letters as if they were distinct. Then divide by 2! to account for 2 duplicate T's, and divide by 2! again for 2 duplicate E's.

More generally...

How many ways can you arrange the letters in “Godoggy”?

$n = 7$ Letters, $k = 4$ Types {G, O, D, Y}

$n_1 = 3, n_2 = 2, n_3 = 1, n_4 = 1$

$$\frac{7!}{3!2!1!1!} = \binom{7}{3,2,1,1}$$

Multinomial coefficients



Multinomial Coefficients

If we have k types of objects (n total), with n_1 of the first type, n_2 of the second, ..., and n_k of the k^{th} , then the number of orderings possible is

$$\binom{n}{n_1, n_2, \dots, n_k} = \frac{n!}{n_1! n_2! \cdots n_k!}$$

Binomial Coefficients – Many interesting and useful properties

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$$\binom{n}{n} = 1$$

$$\binom{n}{1} = n$$

$$\binom{n}{0} = 1$$

Fact. $\binom{n}{k} = \binom{n}{n-k}$

Symmetry in Binomial Coefficients

Fact. $\sum_{k=0}^n \binom{n}{k} = 2^n$

Follows from Binomial Theorem

Fact. $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

Pascal's Identity
(Next lecture)

Binomial Theorem: Idea

$$\begin{aligned}(x + y)^2 &= (x + y)(x + y) \\ &= xx + xy + yx + yy \\ &= x^2 + 2xy + y^2\end{aligned}$$

Poll: What is the coefficient for xy^3 ?

- A. 4
- B. $\binom{4}{1}$
- C. $\binom{4}{3}$
- D. 3

<https://pollev.com/paulbeame028>

$$\begin{aligned}(x + y)^4 &= (x + y)(x + y)(x + y)(x + y) \\ &= xxxx + yyyy + xyxy + yxyy + \dots\end{aligned}$$

Binomial Theorem: Idea

$$(x + y)^n = (x + y) \dots (x + y)$$

Each term is of the form $x^k y^{n-k}$, since each term is made by multiplying exactly n variables, either x or y , one from each copy of $(x + y)$

How many times do we get $x^k y^{n-k}$?

The number of ways to choose x from exactly k of the n copies of $(x + y)$ (the other $n - k$ choices will be y) which is:

$$\binom{n}{k} = \binom{n}{n-k}$$

Binomial Theorem

Theorem. Let $x, y \in \mathbb{R}$ and $n \in \mathbb{N}$ a positive integer. Then,

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

Apply with $x = y = 1$

Many properties of sums of binomial coefficients can be found by plugging in different values of x and y in the Binomial Theorem.

Corollary.

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

Quick Summary

- **k -sequences**: How many length k sequences over alphabet of size n ?
 - Product rule $\rightarrow n^k$
- **k -permutations**: How many length k sequences over alphabet of size n , **without repetition**?
 - Permutation $\rightarrow \frac{n!}{(n-k)!}$
- **k -combinations**: How many size k subsets of a set of size n (**without repetition and without order**)?
 - Combination $\rightarrow \binom{n}{k} = \frac{n!}{k!(n-k)!}$