

CSE 312

Foundations of Computing II

**Lecture 13: Poisson wrap-up
Continuous RV**

Announcements

- PSet 4 due today
- PSet 3 returned yesterday
- Midterm general info is posted on Ed
 - In your section. Closed book . No electronic aids.
- Practice midterm is posted
 - Has format you will see, including 2-page “cheat sheet”.
 - Other practice materials linked also
- Midterm Q&A session next Tuesday 4pm on Zoom

Agenda

- Wrap-up of Poisson RVs ◀
- Continuous Random Variables
- Probability Density Function
- Cumulative Distribution Function

Poisson Random Variables

Definition. A **Poisson random variable** X with parameter $\lambda \geq 0$ is such that for all $i = 0, 1, 2, 3 \dots$,

$$P(X = i) = e^{-\lambda} \cdot \frac{\lambda^i}{i!}$$

General principle:

- Events happen at an average rate of λ per time unit
- Disjoint time intervals independent
- Number of events happening at a time unit X is distributed according to $\text{Poi}(\lambda)$
- Poisson approximates Binomial when n is large, p is small, and np is moderate
- **Sum of independent Poisson is still a Poisson**

Sum of Independent Poisson RVs

Theorem. Let $X \sim \text{Poi}(\lambda_1)$ and $Y \sim \text{Poi}(\lambda_2)$ such that $\lambda = \lambda_1 + \lambda_2$.

Let $Z = X + Y$. For all $z = 0, 1, 2, 3, \dots$,

$$P(Z = z) = e^{-\lambda} \cdot \frac{\lambda^z}{z!}$$

More generally, let $X_1 \sim \text{Poi}(\lambda_1), \dots, X_n \sim \text{Poi}(\lambda_n)$ such that $\lambda = \sum_i \lambda_i$.

Let $Z = \sum_i X_i$

$$P(Z = z) = e^{-\lambda} \cdot \frac{\lambda^z}{z!}$$

Proof

$Z = X + Y$ where $X \sim \text{Poi}(\lambda_1)$ and $Y \sim \text{Poi}(\lambda_2)$ are independent

$$P(Z = z) = \sum_{j=0}^z P(X = j, Y = z - j)$$

Law of total probability

$$= \sum_{j=0}^z P(X = j) P(Y = z - j) = \sum_{j=0}^z e^{-\lambda_1} \cdot \frac{\lambda_1^j}{j!} \cdot e^{-\lambda_2} \cdot \frac{\lambda_2^{z-j}}{(z-j)!} \quad \text{Independence}$$

$$= e^{-\lambda_1 - \lambda_2} \left(\sum_{j=0}^z \frac{1}{j! (z-j)!} \cdot \lambda_1^j \lambda_2^{z-j} \right)$$


$$= e^{-\lambda} \left(\sum_{j=0}^z \frac{z!}{j! (z-j)!} \cdot \lambda_1^j \lambda_2^{z-j} \right) \frac{1}{z!}$$

$$= e^{-\lambda} \cdot (\lambda_1 + \lambda_2)^z \cdot \frac{1}{z!} = e^{-\lambda} \cdot \lambda^z \cdot \frac{1}{z!}$$

Binomial
Theorem

Don't be fooled by this picture: Poisson RVs are discrete

$$\lambda = 5$$
$$p = \frac{5}{n}$$
$$n = 10, 15, 20$$



as $n \rightarrow \infty$, $\text{Binomial}(n, p = \lambda/n) \rightarrow \text{poi}(\lambda)$

Agenda


- Wrap-up of Poisson RVs
- Continuous Random Variables ◀
- Probability Density Function
- Cumulative Distribution Function

Often we want to model experiments where the outcome is not discrete.

Example – Lightning Strike

Lightning strikes a pole within a one-minute time frame


- T = time of lightning strike
- Every time within $[0,1]$ is equally likely
 - Time measured with infinitesimal precision.



The outcome space is not discrete

Lightning strikes a pole within a one-minute time frame


- T = time of lightning strike
- Every point in time within $[0,1]$ is equally likely



Lightning strikes a pole within a one-minute time frame

- T = time of lightning strike
- Every point in time within $[0,1]$ is equally likely


Lightning strikes a pole within a one-minute time frame
• T = time of lightning strike
• Every point in time within $[0,1]$ is equally likely



$$P(0.2 \leq T \leq 0.5) = 0.5 - 0.2 = 0.3$$

Lightning strikes a pole within a one-minute time frame

- T = time of lightning strike
- Every point in time within $[0,1]$ is equally likely




$$P(T = 0.5) = 0$$

Bottom line

- This gives rise to a different type of random variable
- $P(T = x) = 0$ for all $x \in [0,1]$
- Yet, somehow we want
 - $P(T \in [0,1]) = 1$
 - $P(T \in [a, b]) = b - a$
 - ...
- How do we model the behavior of T ?


First try: A discrete approximation

Recall: Cumulative Distribution Function (CDF)




A Discrete Approximation

Probability Mass Function
PMF




Cumulative Distribution Function
CDF



Definition. A **continuous random variable** X is defined by a **probability density function** (PDF) $f_X: \mathbb{R} \rightarrow \mathbb{R}$, such that


Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$



Probability Density Function - Intuition

Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$




Probability Density Function - Intuition


Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$

$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$



Probability Density Function - Intuition



Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$

$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$


$$P(X = y) = P(y \leq X \leq y) = \int_y^y f_X(x) dx = 0$$



Density \neq Probability

$$f_X(y) \neq 0 \quad P(X = y) = 0$$

Probability Density Function - Intuition



Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$


$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$

$$P(X = y) = P(y \leq X \leq y) = \int_y^y f_X(x) dx = 0$$

$$P(X \approx y) \approx P\left(y - \frac{\epsilon}{2} \leq X \leq y + \frac{\epsilon}{2}\right) = \int_{y - \frac{\epsilon}{2}}^{y + \frac{\epsilon}{2}} f_X(x) dx \approx \epsilon f_X(y)$$

What $f_X(x)$ measures: The local **rate** at which probability accumulates

Probability Density Function - Intuition



Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$

$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$

$$P(X = y) = P(y \leq X \leq y) = \int_y^y f_X(x) dx = 0$$

$$P(X \approx y) \approx P\left(y - \frac{\epsilon}{2} \leq X \leq y + \frac{\epsilon}{2}\right) = \int_{y - \frac{\epsilon}{2}}^{y + \frac{\epsilon}{2}} f_X(x) dx \approx \epsilon f_X(y)$$

$$\frac{P(X \approx y)}{P(X \approx z)} \approx \frac{\epsilon f_X(y)}{\epsilon f_X(z)} = \frac{f_X(y)}{f_X(z)} \quad 21$$

Definition. A **continuous random variable** X is defined by a **probability density function** (PDF) $f_X: \mathbb{R} \rightarrow \mathbb{R}$, such that

Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$

$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$

$$P(X = y) = P(y \leq X \leq y) = \int_y^y f_X(x) dx = 0$$

$$P(X \approx y) \approx P\left(y - \frac{\epsilon}{2} \leq X \leq y + \frac{\epsilon}{2}\right) = \int_{y-\frac{\epsilon}{2}}^{y+\frac{\epsilon}{2}} f_X(x) dx \approx \epsilon f_X(y)$$

$$\frac{P(X \approx y)}{P(X \approx z)} \approx \frac{\epsilon f_X(y)}{\epsilon f_X(z)} = \frac{f_X(y)}{f_X(z)}$$




PDF of Uniform RV

$$X \sim \text{Unif}(0,1)$$

Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$

$$f_X(x) = \begin{cases} 1, & x \in [0,1] \\ 0, & x \notin [0,1] \end{cases}$$



$$\int_{-\infty}^{+\infty} f_X(x) dx = \int_0^1 f_X(x) dx = 1 \cdot 1 = 1$$

Probability of Event


$X \sim \text{Unif}(0,1)$

Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$

$$f_X(x) = \begin{cases} 1, & x \in [0,1] \\ 0, & x \notin [0,1] \end{cases}$$

$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$



1. If $0 \leq a$ and $a \leq b \leq 1$

$$P(a \leq X \leq b) = b - a$$

2. If $a < 0$ and $0 \leq b \leq 1$

$$P(a \leq X \leq b) = b$$

3. If $a \geq 0$ and $b > 1$

$$P(a \leq X \leq b) = b - a$$

4. If $a < 0$ and $b > 1$

$$P(a \leq X \leq b) = 1$$

Poll: [pollev/paulbeameo28](#)

A. All of them are correct

B. Only 1, 2, 4 are right

C. Only 1 is right

D. Only 1 and 2 are right

Probability of Event

$X \sim \text{Unif}(0,1)$

Non-negativity: $f_X(x) \geq 0$ for all $x \in \mathbb{R}$

Normalization: $\int_{-\infty}^{+\infty} f_X(x) dx = 1$


$$f_X(x) = \begin{cases} 1, & x \in [0,1] \\ 0, & x \notin [0,1] \end{cases}$$

$$P(a \leq X \leq b) = \int_a^b f_X(x) dx$$

$$P(X = y) = P(y \leq X \leq y) = \int_y^y f_X(x) dx = 0$$

$$P(X \approx y) \approx \epsilon f_X(y) = \epsilon$$

$$\frac{P(X \approx y)}{P(X \approx z)} \approx \frac{\epsilon f_X(y)}{\epsilon f_X(z)} = \frac{f_X(y)}{f_X(z)}$$



PDF of Uniform RV


$$X \sim \text{Unif}(0,0.5)$$



Density \neq Probability

$f_X(x) \gg 1$ is possible!


$$f_X(x) = \begin{cases} 2, & x \in [0,0.5] \\ 0, & x \notin [0,0.5] \end{cases}$$



Uniform Distribution

$X \sim \text{Unif}(a, b)$

$$f_X(x) = \begin{cases} \frac{1}{b-a} & x \in [a, b] \\ 0 & \text{else} \end{cases}$$




$$\int_{-\infty}^{+\infty} f_X(x) dx = (b-a) \frac{1}{b-a} = 1$$

Example. $T \sim \text{Unif}(0,1)$


Probability Density Function

$$f_T(x) = \begin{cases} 1, & x \in [0,1] \\ 0, & x \notin [0,1] \end{cases}$$



Cumulative Distribution Function

$$F_T(x) = P(T \leq x) = \begin{cases} 0 & x \leq 0 \\ ? & 0 \leq x \leq 1 \\ 1 & 1 \leq x \end{cases}$$



Cumulative Distribution Function

Definition. The **cumulative distribution function (cdf)** of X is

$$F_X(a) = P(X \leq a) = \int_{-\infty}^a f_X(x) dx$$

By the fundamental theorem of Calculus $f_X(x) = \frac{d}{dx} F_X(x)$

Therefore: $P(X \in [a, b]) = F_X(b) - F_X(a)$

F_X is monotone increasing, since $f_X(x) \geq 0$. That is $F_X(c) \leq F_X(d)$ for $c \leq d$

$$\lim_{a \rightarrow -\infty} F_X(a) = P(X \leq -\infty) = 0 \quad \lim_{a \rightarrow +\infty} F_X(a) = P(X \leq +\infty) = 1$$

From Discrete to Continuous

	Discrete	Continuous
PMF/PDF	$p_X(x) = P(X = x)$	$f_X(x) \neq P(X = x) = 0$
CDF	$F_X(x) = \sum_{t \leq x} p_X(t)$	$F_X(x) = \int_{-\infty}^x f_X(t) dt$
Normalization	$\sum_x p_X(x) = 1$	$\int_{-\infty}^{\infty} f_X(x) dx = 1$
Expectation	$\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$	$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$